

Truncated Markov bases and Gröbner bases for Integer Programming

Peter N. Malkin

CORE and INMA, Université catholique de Louvain, Belgium
malkin@core.ucl.ac.be

Version of February 2, 2008

Abstract

We present a new algorithm for computing a truncated Markov basis of a lattice. In general, this new algorithm is faster than existing methods. We then extend this new algorithm so that it solves the linear integer feasibility problem with promising results for equality knapsack problems. We also present a novel Gröbner basis approach to solve a particular integer linear program as opposed to previous Gröbner basis methods that effectively solved many different integer linear programs simultaneously. Initial results indicate that this optimisation algorithm performs better than previous Gröbner basis methods.

1 Introduction

Consider the set $\mathcal{F}_{\mathcal{L}}(\nu) := \{x \in \mathbb{N}^n : x \equiv \nu \pmod{\mathcal{L}}\}$ which we call a fiber of the lattice $\mathcal{L}$ where $\nu \in \mathbb{Z}^n$ and $\mathcal{L}$ is a sub-lattice of $\mathbb{Z}^n$. Importantly, the set of feasible solutions to any integer linear program can be equivalently represented in the form $\mathcal{F}_{\mathcal{L}}(\nu')$ for some lattice $\mathcal{L}$ and some $\nu' \in \mathbb{Z}^n$, and conversely, any fiber $\mathcal{F}_{\mathcal{L}}(\nu')$ can be equivalently represented as the set of feasible solutions to an integer linear program (see Appendix A). We use the lattice notation of $\mathcal{F}_{\mathcal{L}}(\nu)$ in this paper since we find it notationally more convenient.

Informally, a Markov basis of the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ is a finite set of vectors in $\mathcal{L}$ such that we can move from any feasible solution to any other feasible solution in a finite number of steps via other feasible solutions using the vectors in the Markov basis. We can step from one feasible solution to another by adding or subtracting a vector in the Markov basis. A Markov basis of the lattice $\mathcal{L}$ is a finite set of vectors that is simultaneously a Markov basis for every possible fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ as ν varies over $\mathbb{Z}^n$. Every lattice has a finite Markov basis, and so, every fiber has a finite Markov basis.

One of the applications of Markov bases is in algebraic statistics to test validity of statistical models via sampling (see Diaconis and Sturmfels [7]). Another area of application is in computational biology for problems arising from phylogenetic trees (see [8]). Markov bases are also needed to perform integer optimisation using Gröbner basis methods. More and more problems are being solved with the growing computational power of computer programs such as **4ti2**, an open source software package for algebraic, geometric, and combinatorial problems on linear spaces ([1]).

In some special situations, finding a Markov basis is straight-forward (see [18]), but in general, this is not the case. There are three main methods for computing a Markov basis of a lattice, (also called a generating set of a lattice): the algorithm of Hosten and Sturmfels in [13] called the “Saturation” algorithm; the algorithm of Bigatti, LaScala, and Robbiano in [3] that we call the “Lift-and-Project” algorithm, and the algorithm of Hemmecke and Malkin in [12] called the “Project-and-Lift” algorithm. Computationally, the Project-and-Lift algorithm is in general the fastest of the three algorithms ([12]).

The above algorithms compute a Markov basis for all fibers of a lattice, but we may only need a Markov basis for one fiber; furthermore, there may be a huge difference between the size of a minimal Markov basis of a lattice and the size of a minimal Markov basis of a fiber. This makes it computationally worthwhile to focus on computing a Markov basis for one fiber. A truncated Markov basis with respect to the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ is a Markov basis of a lattice (all fibers) after removing all vectors that cannot be used to step between two feasible solutions in the fiber; in other words, we remove vectors $u \in \mathcal{L}$ when there do not exist $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ such that $x - y = u$. A vector that cannot step between two feasible solutions in the fiber is never needed in the Markov basis of a fiber, and thus, a truncated Markov basis is thus a Markov basis of the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ but not every fiber. A vector that can step between two feasible solutions in $\mathcal{F}_{\mathcal{L}}(\nu)$ still may not be necessary in a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$, but hopefully, there are not too many unnecessary vectors in the truncated Markov basis.

The new algorithm for computing truncated Markov bases presented in this paper in Section 5 is based upon the Project-and-Lift algorithm combined with a truncated Gröbner basis algorithm (see [18]), which is described in Section 4. Previously, there were two known methods. The first straight-forward method computes a Markov basis of a lattice and then truncates the Markov basis. The second method computes a truncated Graver basis of the lattice ([11]), which is a superset of a truncated Gröbner basis; however, a truncated Graver basis might be a lot larger than a minimal truncated Markov basis. The new method is in general much faster than these two methods.

In Section 6, we present a new algorithm to compute a feasible solution of a fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ (i.e. find a point $x \in \mathcal{F}_{\mathcal{L}}(\nu)$ if one exists) by using an extension of the truncated Markov basis algorithm. At the same time as finding a Markov basis, the truncated Markov basis algorithm can also compute a feasible solution without much additional computation. The notions of Markov bases and feasibility are indeed strongly related. Recall that finding a feasible solution of a fiber is equivalent to finding a feasible solution of a linear integer program. Feasible solutions are not only interesting in themselves, but are also needed to perform integer programming optimisation using a Gröbner basis (see below). We also show how to compute a feasible solution of a finite set of different fibers $\mathcal{F}_{\mathcal{L}}(\nu)$ simultaneously. This is also extended to the truncated case where we focus on one particular fiber. We present the promising results of applying the algorithm to solve equality constrained integer knapsacks ([2]).

In Section 7, we solve linear integer programs using Gröbner bases, which in this context are also known as *test sets* for integer linear programs. Test sets were first introduced by Graver in [10] (see for example [22]). A Gröbner basis of a fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ with respect to some term order $\succ$ is a finite set of vectors in $\mathcal{L}$ such that we can improve every non $\succ$ -minimal point $x \in \mathcal{F}_{\mathcal{L}}(\nu)$ by subtracting a vector in the Gröbner basis. A Gröbner basis is thus a set of *augmenting* or *improving* vectors. So, using a $\succ$ -Gröbner basis, given some initial feasible solution of the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$, we can find the $\succ$ -minimal point in the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ by iteratively improving the feasible solution until we can no longer do so, in which case, we must have a $\succ$ -minimal solution. We call the problem of finding the $\succ$ -minimal point in a fiber a lattice program; that is, the problem

$IP_{\mathcal{L}, \succ}(\nu) := \min_{\succ} \{x : x \in \mathcal{F}_{\mathcal{L}}(\nu)\}$. A Gröbner basis of a lattice is a finite set of vectors in $\mathcal{L}$ that is simultaneously a Gröbner basis for every fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ as ν varies over $\mathbb{Z}^n$. Any integer linear program can be expressed as a lattice program and conversely any lattice program may be written as an integer linear program, and thus, the concepts of integer linear programs and lattice programs are equivalent (see Appendix A).

The first Gröbner basis methods for solving a linear integer program constructed Gröbner bases of lattices and thus effectively solved the program for all fibers simultaneously ([5]). This makes the approach appealing if we wish to solve the program for many different fibers; however, if we wish to solve the program for only one fiber then this is a disadvantage because we often do much more work than necessary. By using the structure of a fiber, the new algorithm presented here focuses on solving the program for a specific fiber. To achieve this, the algorithm does two things: it solves a hierarchy of group relaxations to avoid computing with unnecessary constraints and it also applies truncation methods which are strengthened by using the cost function. Solving a hierarchy of group relaxations was explored in [9, 23, 20, 17], and truncated methods were explored in [18]; however, combining these two approaches and strengthening truncation by using the cost function has not been done before.

Gröbner bases and Markov bases have corresponding concepts in computational algebraic geometry (see [5]) although we do not present it here. Instead, the approach in this paper follows the geometric approach in [12, 18, 19, 21, 22].

All computations in the paper were done using `4ti2` version 1.3 on a Intel Pentium4 3.0GHz machine running Linux. All timings given are rounded to the nearest one hundredth of a second.

2 Truncated Markov Bases

In this section, we define Markov bases of fibers, Markov bases of lattices, and Markov bases of Gröbner bases. This notion of truncation has been explored in [21] and [18], but only for computing truncated Gröbner bases, and we apply it here to Markov bases.

A **lattice** is a set $\mathcal{L} \subseteq \mathbb{Z}^n$ where $\mathcal{L} = \mathbb{Z}(S) = \{\sum_{i=1}^k \lambda_i s^i : \lambda \in \mathbb{Z}^k\}$ for some finite set $S = \{s^1, \dots, s^k\} \subseteq \mathbb{Z}^n$. If $\mathcal{L} = \mathbb{Z}(S)$, then we say that S spans $\mathcal{L}$, and if S is inclusion-minimal, then we call S a basis of $\mathcal{L}$.

Given a lattice $\mathcal{L} \subseteq \mathbb{Z}^n$, and a vector $\nu \in \mathbb{Z}^n$, we define the set

$$\mathcal{F}_{\mathcal{L}}(\nu) := \{x \in \mathbb{N}^n : x \equiv \nu \pmod{\mathcal{L}}\} = \{x \in \mathbb{N}^n : x - \nu \in \mathcal{L}\}$$

that we call a **fiber** of the lattice $\mathcal{L}$.

Given a lattice $\mathcal{L} \subseteq \mathbb{Z}^n$, a vector $\nu \in \mathbb{Z}^n$, and a set $S \subseteq \mathcal{L}$, we define the *fiber graph* $\mathcal{G}_{\mathcal{L}}(\nu, S)$ to be the *undirected* graph with nodes $\mathcal{F}_{\mathcal{L}}(\nu)$ and edges (x, y) if $x - y \in S$ or $y - x \in S$ for $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$.

Definition 2.1. *Given $\nu \in \mathbb{Z}^n$, we call a set $S \subseteq \mathcal{L}$ a **Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$** if the graph $\mathcal{G}_{\mathcal{L}}(\nu, S)$ is connected. The set S is called a **Markov basis of $\mathcal{L}$** if it is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ for every $\nu \in \mathbb{Z}^n$.*

We remind the reader that connectedness of $\mathcal{G}_{\mathcal{L}}(\nu, S)$ simply states that between each pair $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ there exists a path from x to y in the graph $\mathcal{G}_{\mathcal{L}}(\nu, S)$.

Example 2.2. Let $S := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1)\}$, and let $\mathcal{L} \subseteq \mathbb{Z}^6$ be the lattice spanned by S . By definition, S is a spanning set of $\mathcal{L}$, but S is not a Markov basis of $\mathcal{L}$. Observe that $\mathcal{L} = \mathcal{L}_A := \{u : Au = \mathbf{0}, u \in \mathbb{Z}^n\}$ where

$$A = (\tilde{A}, I), \quad \tilde{A} = \begin{pmatrix} -2 & -3 \\ +2 & -1 \\ +1 & +2 \\ -1 & +1 \end{pmatrix}, \quad \text{and } I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

So, for every $\nu \in \mathbb{Z}^6$, $\mathcal{F}_{\mathcal{L}}(\nu) = F_A(b) = \{(x, s) : \tilde{A}x + Is = b, x \in \mathbb{N}^2, s \in \mathbb{N}^4\}$ where $b = A\nu \in \mathbb{Z}^4$. Hence, the projection of $\mathcal{F}_{\mathcal{L}}(\nu)$ onto the (x_1, x_2) -plane is the set of integer points in the polyhedron $\{x \in \mathbb{R}_+^n : \tilde{A}x \leq b\}$, and the s variables are the slack variables. Consider $\nu := (2, 2, 4, 2, 5, 1)$; then, $\mathcal{F}_{\mathcal{L}}(\nu) = F_A(b)$ where $b = A\nu = (-6, 4, 11, 1)$ (see Figure 1a).




Figure 1: The set $\mathcal{F}_{\mathcal{L}}(\nu)$ and the graphs $\mathcal{G}_{\mathcal{L}}(\nu, S)$ and $\mathcal{G}_{\mathcal{L}}(\nu, S')$ projected onto the (x_1, x_2) -plane.

The graph $\mathcal{G}_{\mathcal{L}}(\nu, S)$ is not connected because the point $(3, 4, 12, 2, 0, 0) \in \mathcal{F}_{\mathcal{L}}(\nu)$ is disconnected (see Figure 1b). Let $S' := S \cup \{(1, 1, 5, -1, -3, 0)\}$. The graph of $\mathcal{G}_{\mathcal{L}}(\nu, S')$ is now connected (see Figure 1c). Thus, S' is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$.

A **truncated Markov basis** is a special type of Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ for some $\nu \in \mathbb{Z}^n$ that is not necessarily a Markov basis of all fibers. Essentially, a truncated Markov basis with respect to the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ is a Markov basis of $\mathcal{L}$ after removing all vectors $u \in \mathcal{L}$ for which there does not exist $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ such that $x - y = u$. We call the act of removing such vectors *truncation*. More formally, let G be a Markov basis of $\mathcal{L}$; then, the set $S := \{u \in G : u = x - y \text{ for some } x, y \in \mathcal{F}_{\mathcal{L}}(\nu)\}$ is a truncated Markov basis. Any vector $u \in \mathcal{L}$ for which there does not exist $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ such that $u = x - y$ can never be an edge in a fiber graph. Hence, we never need such a vector u in a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$. Therefore, S must be a Markov basis of the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$.

The set S above is also a Markov basis of other related fibers. Let $\nu' \in \mathbb{Z}^n$ where $\mathcal{F}_{\mathcal{L}}(\nu') \neq \emptyset$ and $\mathcal{F}_{\mathcal{L}}(\nu - \nu') \neq \emptyset$. The set S is also a Markov basis of the fiber $\mathcal{F}_{\mathcal{L}}(\nu')$. Let $u \in \mathcal{L}$ for which there exists $x, y \in \mathcal{F}_{\mathcal{L}}(\nu')$ such that $u = x - y$, and let $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')$. Then, $x + \gamma, y + \gamma \in \mathcal{F}_{\mathcal{L}}(\nu)$, and moreover, $u = (x + \gamma) - (y + \gamma)$; thus, u would not be removed during truncation. So, any vector needed in a Markov basis of the fiber $\mathcal{F}_{\mathcal{L}}(\nu')$ would not be removed by truncation, and therefore, S is still a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu')$. The set S is thus a Markov basis of the following set of fibers:

$$\mathcal{B}_{\mathcal{L}}(\nu) := \{\nu' \in \mathbb{Z}^n : \mathcal{F}_{\mathcal{L}}(\nu') \neq \emptyset \text{ and } \mathcal{F}_{\mathcal{L}}(\nu - \nu') \neq \emptyset\}.$$

This property of a truncated Markov basis is the defining property of truncated Markov bases.

Definition 2.3. Given $\nu \in \mathbb{Z}^n$, we call a set $S \subseteq \mathcal{L}$ a ν -truncated Markov basis of $\mathcal{L}$ if S is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu')$ for every $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$.

Note that if $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$, then $\nu \in \mathcal{B}_{\mathcal{L}}(\nu)$ since $\mathcal{F}_{\mathcal{L}}(\mathbf{0}) \neq \emptyset$. Therefore, a ν -truncated Markov basis is by definition a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$, but a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ is not necessarily a ν -truncated Markov basis. Moreover, a ν -truncated Markov basis of $\mathcal{L}$ is not necessarily a Markov basis of $\mathcal{L}$. In the special case where $\mathcal{F}_{\mathcal{L}}(\nu) = \emptyset$, we have $\mathcal{B}_{\mathcal{L}}(\nu) = \emptyset$, which is consistent since by definition an empty set is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ if $\mathcal{F}_{\mathcal{L}}(\nu) = \emptyset$.

Additionally, given a vector $u \in \mathcal{L}$, there exists $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ where $x - y = u$ if and only if $\mathcal{F}_{\mathcal{L}}(\nu - u^+) \neq \emptyset$ since $x - y = u$ means that $x = \gamma + u^+$ and $y = \gamma + u^-$ for some $\gamma \in \mathbb{N}^n$ in which case $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu - u^+) \neq \emptyset$. Here, $u^+ \in \mathbb{N}^n$ is the positive part of u and $u^- \in \mathbb{N}^n$ is the negative part, or in other words, $u_i^+ = \max\{0, u_i\}$ and $u_i^- = \max\{0, -u_i\}$ for all $i = 1, \dots, n$. Moreover, since $u^+ \in \mathcal{F}_{\mathcal{L}}(u^+) \neq \emptyset$, we have that there exists $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ where $x - y = u$ if and only if $u^+ \in \mathcal{B}_{\mathcal{L}}(\nu)$.

The set $\mathcal{B}_{\mathcal{L}}(\nu)$ has some interesting properties. Given a set $S \subseteq \mathcal{L}$, the connectivity of the graph $\mathcal{G}_{\mathcal{L}}(\nu', S)$ for the fibers in $\mathcal{B}_{\mathcal{L}}(\nu)$ is strongly related to the connectivity of $\mathcal{G}_{\mathcal{L}}(\nu, S)$. Note that for $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$, we have $\gamma + \mathcal{F}_{\mathcal{L}}(\nu') \subseteq \mathcal{F}_{\mathcal{L}}(\nu)$ for every $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')$ where $\gamma + \mathcal{F}_{\mathcal{L}}(\nu') = \{\gamma + x : x \in \mathcal{F}_{\mathcal{L}}(\nu')\}$. So, given $S \subseteq \mathcal{L}$, any path $(x^0, \dots, x^k)$ in $\mathcal{G}_{\mathcal{L}}(\nu', S)$ can be translated by γ to a path $(x^0 + \gamma, \dots, x^k + \gamma)$ in $\mathcal{G}_{\mathcal{L}}(\nu, S)$ for every $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')$. Hence, if S is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu')$, then any two points in $\gamma + \mathcal{F}_{\mathcal{L}}(\nu') \subseteq \mathcal{F}_{\mathcal{L}}(\nu)$ are connected in $\mathcal{G}_{\mathcal{L}}(\nu, S)$. Moreover, we have $\mathcal{F}_{\mathcal{L}}(\nu') + \mathcal{F}_{\mathcal{L}}(\nu - \nu') \subseteq \mathcal{F}_{\mathcal{L}}(\nu)$ where $\mathcal{F}_{\mathcal{L}}(\nu') + \mathcal{F}_{\mathcal{L}}(\nu - \nu') = \{x + y : x \in \mathcal{F}_{\mathcal{L}}(\nu'), y \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')\}$. Note that $\nu - \nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$ when $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$. If S is both a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu')$ and a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu - \nu')$, then any two points in $\mathcal{F}_{\mathcal{L}}(\nu') + \mathcal{F}_{\mathcal{L}}(\nu - \nu')$ are connected in $\mathcal{G}_{\mathcal{L}}(\nu, S)$. This is shown as follows. Any two points in $\mathcal{F}_{\mathcal{L}}(\nu') + \mathcal{F}_{\mathcal{L}}(\nu - \nu')$ can be written in the form $x^1 + y^1$ and $x^2 + y^2$ where $x^1, x^2 \in \mathcal{F}_{\mathcal{L}}(\nu')$ and $y^1, y^2 \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')$. Now, from above, the points $x^1 + y^1, x^2 + y^1 \in y^1 + \mathcal{F}_{\mathcal{L}}(\nu')$ are connected in $\mathcal{G}_{\mathcal{L}}(\nu, S)$ and the points $x^2 + y^1, x^2 + y^2 \in x^2 + \mathcal{F}_{\mathcal{L}}(\nu - \nu')$ are connected in $\mathcal{G}_{\mathcal{L}}(\nu, S)$; hence, the points $x^1 + y^1$ and $x^2 + y^2$ are connected in $\mathcal{G}_{\mathcal{L}}(\nu, S)$ as required.

Example 2.4. Consider again the set $S' := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1), (1, 1, 5, -1, -3, 0)\}$ and the lattice $\mathcal{L}$ from above. We saw previously that S' is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ where $\nu = (2, 2, 4, 2, 5, 1)$; however, S' is not a ν -truncated Markov basis of $\mathcal{L}$. Consider $\nu' = (2, 2, 4, 2, 0, 0)$ (see Figure 2a). Note that $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$ since $\nu - \nu' = (0, 0, 0, 0, 5, 1) \in \mathcal{F}_{\mathcal{L}}(\nu - \nu') \neq \emptyset$.




Figure 2: The sets $\mathcal{F}_{\mathcal{L}}(\nu')$, $\mathcal{F}_{\mathcal{L}}(\nu'')$, and $\mathcal{F}_{\mathcal{L}}(\nu - \nu'')$ projected onto the (x_1, x_2) -plane.

The graph $\mathcal{G}_{\mathcal{L}}(\nu', S')$ is disconnected since there are only two feasible points in $\mathcal{F}_{\mathcal{L}}(\nu') = \{(2, 1, 1, 1, 2, 1), (2, 2, 4, 2, 0, 0)\}$. The vector $(0, 1, 3, 1, -2, -1)$ gives the unique minimal Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu')$, and

hence, it must be in a ν -truncated Markov basis of $\mathcal{L}$. The set $S'' = S' \cup (0, 1, 3, 1, -2, -1)$ is a ν -truncated Markov basis of $\mathcal{L}$.

Although S'' is a ν -truncated Markov basis of $\mathcal{L}$, it is not a Markov basis for every $\nu'' \in \mathbb{Z}^n$. Consider $\nu'' = (0, 1, 0, 5, 0, 0)$. The graph $\mathcal{G}_{\mathcal{L}}(\nu'', S'')$ is disconnected since there are only two feasible points in $\mathcal{F}_{\mathcal{L}}(\nu'') = \{(2, 0, 1, 0, 0, 3), (0, 1, 0, 5, 0, 0)\}$ (see Figure 2b). Observe that $\nu'' \notin \mathcal{B}_{\mathcal{L}}(\nu)$ since $\mathcal{F}_{\mathcal{L}}(\nu - \nu'') = \emptyset$ (see Figure 2c).

In some situations, the ν -truncated Markov basis of $\mathcal{L}$ may be empty: a minimal ν -truncated Markov basis is empty if and only if $|\mathcal{F}_{\mathcal{L}}(\nu)| \leq 1$ since then the fiber has zero or one elements. On the other hand, a minimal ν -truncated Markov basis of $\mathcal{L}$ may also be a Markov basis of $\mathcal{L}$. So, the degree to which truncation affects the size of the Markov basis varies between the two extremes of an empty set and a Markov basis of a lattice.

3 Truncated Gröbner bases

In this section, we define Gröbner bases of fibers, Gröbner bases of lattices, and truncated Gröbner bases (see [21, 18]).

First, we need to define term orders. We call $\succ$ a **term ordering** for $\mathcal{L}$ if

- (i). $\succ$ is a total ordering on the set $\mathcal{F}_{\mathcal{L}}(\nu)$ for every $\nu \in \mathbb{Z}^n$,
- (ii). there is a unique $\succ$ -minimal solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ for every $\nu \in \mathbb{Z}^n$ for which $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$, and
- (iii). $\succ$ is an additive ordering meaning that for all $\nu \in \mathbb{Z}^n$ and for all $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$, if $x \succ y$, then $x + \gamma \succ y + \gamma$ for every $\gamma \in \mathbb{N}^n$ (note that $x + \gamma, y + \gamma \in \mathcal{F}_{\mathcal{L}}(\nu + \gamma)$).

The most common term orders are the lexicographic term ordering and the degree reverse lexicographic term ordering (see for example [6]).

Definition 3.1. Given $\nu \in \mathbb{Z}^n$ and a term order $\succ$, we call $G \subseteq \mathcal{L}$ a **$\succ$ -Gröbner basis** of $\mathcal{F}_{\mathcal{L}}(\nu)$ if for every $x \in \mathcal{F}_{\mathcal{L}}(\nu)$, either x is the unique $\succ$ -minimal element of $\mathcal{F}_{\mathcal{L}}(\nu)$ or there exists a vector $u \in G$ such that $x - u \in \mathcal{F}_{\mathcal{L}}(\nu)$ and $x \succ x - u$. The set G is called a **$\succ$ -Gröbner basis of $\mathcal{L}$** if it is a Gröbner basis for every $\nu \in \mathbb{Z}^n$.

Analogously to truncated Markov bases, we define truncated Gröbner bases.

Definition 3.2. Given $\nu \in \mathbb{Z}^n$, we call a set $G \subseteq \mathcal{L}$ a **ν -truncated $\succ$ -Gröbner basis of $\mathcal{L}$** if G is a Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu')$ for every $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$.

As in the Markov basis case, a ν -truncated $\succ$ -Gröbner basis is by definition a $\succ$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu)$, but a $\succ$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ is not necessarily a ν -truncated $\succ$ -Gröbner basis and furthermore, a ν -truncated $\succ$ -Gröbner basis of $\mathcal{L}$ is not necessarily a $\succ$ -Gröbner basis of $\mathcal{L}$.

We can solve *lattice programs* using Gröbner bases. Given a lattice $\mathcal{L}$, a vector ν , and a term order $\succ$, the problem

$$IP_{\mathcal{L}, \succ}(\nu) := \min_{\succ} \{x : x \in \mathcal{F}_{\mathcal{L}}(\nu)\}$$

is called a **lattice program**. Given some initial feasible solution $x \in \mathcal{F}_{\mathcal{L}}(\nu)$, we can solve the lattice program $IP_{\mathcal{L},\succ}(\nu)$ using a Gröbner basis G of the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ by iteratively improving the feasible solution using vectors in G . This process constructs a $\succ$ -decreasing path in the graph $\mathcal{G}_{\mathcal{L}}(\nu, G)$ from the initial feasible solution x to the unique $\succ$ -minimal solution where a path $(x^0, \dots, x^k)$ in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ is **$\succ$ -decreasing** if $x^i \succ x^{i+1}$ for $i = 0, \dots, k-1$. This gives us an equivalent way of defining Gröbner bases in terms of paths in the graph $\mathcal{G}_{\mathcal{L}}(\nu, G)$ in Lemma 3.3 below. Firstly, note that for a vector $u \in \mathcal{L}$, where $x, x-u \in \mathcal{F}_{\mathcal{L}}(\nu)$ and $x \succ x-u$, we must have $u^+ \succ u^-$. This property follows since $\succ$ is an additive ordering. Thus, we only need consider vectors in the set $\mathcal{L}_{\succ} := \{u \in \mathcal{L} : u^+ \succ u^-\}$.

Lemma 3.3. *Given $\nu \in \mathbb{Z}^n$, $G \subseteq \mathcal{L}_{\succ}$ is a $\succ$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ if and only if for every $x \in \mathcal{F}_{\mathcal{L}}(\nu)$ there exists a decreasing path in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ from x to the unique $\succ$ -minimal element in $\mathcal{F}_{\mathcal{L}}(\nu)$.*

Importantly, if $G \subseteq \mathcal{L}_{\succ}$ is a $\succ$ -Gröbner basis, then G is a Markov basis of $\mathcal{L}$ since given $x, y \in \mathcal{G}_{\mathcal{L}}(\nu, G)$ for some $\nu \in \mathbb{Z}^n$, there exists a $\succ$ -decreasing path from x to the unique $\succ$ -minimal element in $\mathcal{F}_{\mathcal{L}}(\nu)$ and from y to the same element, and thus, x and y are connected in $\mathcal{G}_{\mathcal{L}}(\nu, G)$.

We can use Gröbner bases to solve the integer program

$$IP_{\mathcal{L},c}(\nu) := \min\{cx : x \in \mathcal{F}_{\mathcal{L}}(\nu)\}$$

given a lattice $\mathcal{L}$, a vector $\nu \in \mathbb{Z}^n$, and a cost vector $c \in \mathbb{Z}^n$. To solve $IP_{\mathcal{L},c}(\nu)$, we solve instead a lattice program $IP_{\mathcal{L},\succ}(\nu)$ for some *term order* $\succ$ that is compatible with c .

Given a vector $c \in \mathbb{Z}^n$, we say that a vector a term order $\succ$ is **compatible** with c if the optimal solution of $IP_{\mathcal{L},\succ}(\nu)$ is also an optimal solution of $IP_{\mathcal{L},c}(\nu)$ for all $\nu \in \mathbb{Z}^n$ where $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$. We can easily construct a c compatible order $\succ_c$ given some (tie-breaking) term ordering $\succ$ as follows: $x \succ_c y$ if $cx > cy$, or $cx = cy$ and $x \succ y$. We must be a little careful here though since $\succ_c$ is not necessarily a term order. The ordering $\succ_c$ satisfies conditions (i) and (iii) for being a term order, but condition (ii) is not always satisfied. The ordering $\succ_c$ is a term ordering if and only if $IP_{\mathcal{L},c}(\nu)$ has an optimal solution for every $\nu \in \mathbb{Z}^n$ where $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$. Note that $IP_{\mathcal{L},c}(\nu)$ has an optimal solution for every $\nu \in \mathbb{Z}^n$ where $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$ if and only if $IP_{\mathcal{L},c}(\mathbf{0}) := \min\{cx : x \in \mathcal{L}, x \in \mathbb{N}^n\} = 0$, and we can check whether $IP_{\mathcal{L},c}(\mathbf{0}) = 0$ using linear programming.

Conversely, given any term order $\succ$ there always exists a compatible $c \in \mathbb{Z}^n$ such that $IP_{\mathcal{L},c}(\nu)$ has a unique optimal solution that is also the optimal solution of $IP_{\mathcal{L},\succ}(\nu)$ for all $\nu \in \mathbb{Z}^n$ where $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$ ([16]). So, solving integer programs $IP_{\mathcal{L},c}(\nu)$ is essentially equivalent to solving lattice programs.

4 Computing truncated Gröbner bases

In this section, we describe how to compute truncated Gröbner bases. We first describe how to compute truncated Markov bases since the algorithm for computing truncated Gröbner bases is used to compute truncated Markov bases. We present existing algorithms for truncated Gröbner bases including a new approaches to truncation. The structure of this section follows closely from [12] except that we now deal with truncation, and so, we have omitted the proofs of those results in this section that correspond closely to results in [12].

We now describe Gröbner bases in terms of reduction paths, so that we avoid explicitly mentioning the $\succ$ -minimal solution of every fiber. A path $(x^0, \dots, x^k)$ in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ is a $\succ$ -**reduction path** if for all $i \in \{1, \dots, k-1\}$, we have either $x^0 \succ x^i$ or $x^k \succ x^i$. For example, see Figure 3.




Figure 3: Reduction path between x and y .

Lemma 4.1. *Given $\nu \in \mathbb{Z}^n$, a set $G \subseteq \mathcal{L}_{\succ}$ is a $\succ$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ if and only if for each pair $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$, there exists a $\succ$ -reduction path in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ between x and y .*

Checking for a given $G \subseteq \mathcal{L}_{\succ}$ whether there exists a $\succ$ -reduction path in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ for every $\nu \in \mathbb{Z}^n$ and for each pair $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ involves many situations that need to be checked. In fact, far fewer checks are needed: we only need to check for a $\succ$ -reduction path from x to y if there exists a $\succ$ -**critical path** from x to y .

Definition 4.2. *Given $G \subseteq \mathcal{L}_{\succ}$ and $\nu \in \mathbb{Z}^n$, a path (x, z, y) in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ is a $\succ$ -**critical path** if $z \succ x$ and $z \succ y$.*

If (x, z, y) is a $\succ$ -critical path in $\mathcal{G}_{\mathcal{L}}(\nu, G)$, then $x + u = z = y + v$ for some pair $u, v \in G$, in which case, we call (x, z, y) a $\succ$ -critical path for (u, v) (see Figure 4).




Figure 4: A critical path for (u, v) between x, z , and y .

Lemma 4.3. *Given $\nu \in \mathbb{Z}^n$, let $G \subseteq \mathcal{L}_{\succ}$ where G is a Markov basis of $\mathcal{F}_{\mathcal{L}}(\nu)$. G is a $\succ$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu)$ if and only if there exists a $\succ$ -reduction path between x' and y' for every $\succ$ -critical path (x', z', y') in $\mathcal{G}_{\mathcal{L}}(\nu, G)$.*

We can extend Lemma 4.3 to ν -truncated Gröbner bases. It is a straight-forward consequence of Lemma 4.3, but nevertheless, it is worthwhile stating explicitly.

Lemma 4.4. *Given $\nu \in \mathbb{Z}^n$, let $G \subseteq \mathcal{L}_{\succ}$ where G is a ν -truncated Markov basis of $\mathcal{L}$. G is a ν -truncated $\succ$ -Gröbner basis of $\mathcal{L}$ if and only if there exists a $\succ$ -reduction path between x' and y' for every $\succ$ -critical path (x', z', y') in $\mathcal{G}_{\mathcal{L}}(\nu', G)$ for all $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$.*

It is not necessary to check for a $\succ$ -reduction path from x' to y' for every $\succ$ -critical path (x', y', z') in $\mathcal{G}_{\mathcal{L}}(\nu', G)$ for all $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$. Consider the case where there exists another $\succ$ -critical path (x'', y'', z'') in $\mathcal{G}_{\mathcal{L}}(\nu'', G)$ for some $\nu'' \in \mathbb{Z}^n$ such that $(x', y', z') = (x'' + \gamma, y'' + \gamma, z'' + \gamma)$ for some

$\gamma \in \mathbb{N}^n$. Then, a $\succ$ -reduction path from x'' to y'' in $\mathcal{G}_{\mathcal{L}}(\nu'', G)$ translates by γ to a $\succ$ -reduction path from x' to y' in $\mathcal{G}_{\mathcal{L}}(\nu', G)$. Moreover, $\nu'' \in \mathcal{B}_{\mathcal{L}}(\nu)$, since $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu' - \nu'') \neq \emptyset$ which implies that $\mathcal{F}_{\mathcal{L}}(\nu - \nu'') \neq \emptyset$ because $\mathcal{F}_{\mathcal{L}}(\nu - \nu') \neq \emptyset$. Thus, we only need to check for a $\succ$ -reduction path from x'' to y'' .

We call a $\succ$ -critical path (x, y, z) **minimal** if there does not exist another $\succ$ -critical path (x', y', z') such that $(x, y, z) = (x' + \gamma, y' + \gamma, z' + \gamma)$ for some $\gamma \in \mathbb{N}^n$ where $\gamma \neq 0$, or equivalently, $\min\{x_i, y_i, z_i\} = 0$ for all $i = 1, \dots, n$. Consequently, if there exists a $\succ$ -reduction path between x' and y' for all minimal $\succ$ -critical paths (x', y', z') in $\mathcal{G}_{\mathcal{L}}(\nu', G)$ for some $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$, then there exists a $\succ$ -reduction path between x'' and y'' for all $\succ$ -critical paths (x'', y'', z'') in $\mathcal{G}_{\mathcal{L}}(\nu'', G)$ for some $\nu'' \in \mathcal{B}_{\mathcal{L}}(\nu)$. Also, for each pair of vectors $u, v \in G$, there exists a unique minimal $\succ$ -critical path $(x^{(u,v)}, z^{(u,v)}, y^{(u,v)})$ in $\mathcal{G}_{\mathcal{L}}(\nu^{(u,v)}, G)$ determined by $z^{(u,v)} := \max\{u^+, v^+\}$ component-wise, $x^{(u,v)} := z^{(u,v)} - u$, $y^{(u,v)} := z^{(u,v)} - v$, and $\nu^{(u,v)} := z^{(u,v)}$. So, any other $\succ$ -critical path for (u, v) is of the form $(x^{(u,v)} + \gamma, z^{(u,v)} + \gamma, y^{(u,v)} + \gamma)$ for some $\gamma \in \mathbb{N}^n$. Using minimal $\succ$ -critical paths, we can rewrite Lemma 4.4, so that we only need to check for a *finite* number of $\succ$ -reduction paths.

Lemma 4.5. *Let $G \subseteq \mathcal{L}_{\succ}$ and $\nu \in \mathbb{Z}^n$. The set G is a ν -truncated $\succ$ -Gröbner basis of $\mathcal{L}$ if and only if for all $u, v \in G$ where $\nu^{(u,v)} \in \mathcal{B}_{\mathcal{L}}(\nu)$, there exists a $\succ$ -reduction path between $x^{(u,v)}$ and $y^{(u,v)}$ in $\mathcal{G}_{\mathcal{L}}(\nu^{(u,v)}, G)$.*

We now turn Lemma 4.5 into an algorithmic tool. Algorithm 2 below, called a completion procedure ([4]), starts from a ν -truncated Markov basis and computes a ν -truncated $\succ$ -Gröbner basis. An important part of this algorithm is checking whether $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$ for some $\nu' \in \mathbb{Z}^n$. How exactly we perform this check in practice is discussed at length after first presenting the overall algorithm.

Given a set $S \subseteq \mathcal{L}$, the completion procedure first sets $G := S$, and then directs all vectors in G according to $\succ$ such that $G \subseteq \mathcal{L}_{\succ}$. It also removes from the set G any vectors $u \in G$ such that $u^+ \notin \mathcal{B}_{\mathcal{L}}(\nu)$ – recall that these vectors are not needed in a truncated Markov basis. Note that at this point $\mathcal{G}_{\mathcal{L}}(\nu', S) = \mathcal{G}_{\mathcal{L}}(\nu', G)$ for all $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$, and thus, G is also a ν -truncated Markov basis of $\mathcal{L}$. The completion procedure then determines whether the set G satisfies Lemma 4.5; in other words, it tries to find a reduction path from $x^{(u,v)}$ to $y^{(u,v)}$ for every pair $u, v \in G$ where $\nu^{(u,v)} \in \mathcal{B}_{\mathcal{L}}(\nu)$. If G satisfies Lemma 4.5, then we are done. Otherwise, no $\succ$ -reduction path was found for some (u, v) , in which case, we add a vector to G so that a $\succ$ -reduction path exists, and then again, test whether G satisfies Lemma 4.5 and so on.

To check for a $\succ$ -reduction path from $x^{(u,v)}$ to $y^{(u,v)}$, we construct a maximal $\succ$ -decreasing path in $\mathcal{G}_{\mathcal{L}}(\nu^{(u,v)}, G)$ from $x^{(u,v)}$ to some x' and from $y^{(u,v)}$ to some y' using the “Normal Form Algorithm” (Algorithm 1 below). If $x' = y'$, then we have found a $\succ$ -reduction path from $x^{(u,v)}$ to $y^{(u,v)}$. Otherwise, we add the vector $r \in \mathcal{L}_{\succ}$ to G where $r := x' - y'$ if $x' \succ y'$, and $r := y' - x'$ otherwise, and then, there is now a $\succ$ -reduction path from $x^{(u,v)}$ to $y^{(u,v)}$ in $\mathcal{G}_{\mathcal{L}}(\nu^{(u,v)}, G)$.

Algorithm 1 Normal Form Algorithm

Input: a vector $x \in \mathbb{N}^n$ and a set $G \subseteq \mathcal{L}_{\succ}$.

Output: a vector x' where there is a maximal $\succ$ -decreasing path from x to x' in $\mathcal{G}_{\mathcal{L}}(x, G)$.

```

 $x' := x$ 
while there is exists  $u \in G$  such that  $u^+ \leq x'$  do
     $x' := x' - u$ 
end while
return  $x'$ 

```

Algorithm 2 Truncated completion procedure

Input: a vector $\nu \in \mathbb{Z}^n$, a term ordering $\succ$, and a ν -truncated Markov basis $S \subseteq \mathcal{L}$.

Output: a ν -truncated Gröbner basis $G \subseteq \mathcal{L}_\succ$.

$G := \{u : u^+ \succ u^-, u \in S\} \cup \{-u : u^- \succ u^+, u \in S\}$

$G := \{u : u \in G, u^+ \in \mathcal{B}_\mathcal{L}(\nu)\}$

$C := \{(u, v) : u, v \in G, \nu^{(u,v)} \in \mathcal{B}_\mathcal{L}(\nu)\}$

while $C \neq \emptyset$ **do**

 Select $(u, v) \in C$

$C := C \setminus \{(u, v)\}$

$r := \mathcal{NF}(x^{(u,v)}, G) - \mathcal{NF}(y^{(u,v)}, G)$

if $r \neq 0$ **then**

if $r^- \succ r^+$ **then** $r := -r$

$C := C \cup \{(r, s) : s \in G, \nu^{(r,s)} \in \mathcal{B}_\mathcal{L}(\nu)\}$

$G := G \cup \{r\}$

end if

end while

return G .

We write $\mathcal{NF}(x, G)$ for the output of the Normal Form Algorithm and we write $\mathcal{CP}_\nu(\succ, S)$ for the output of the Completion Procedure.

There is a trade off between the computational benefit of computing a ν -truncated Markov basis (computing a smaller set) and the computational cost of computing whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$ for some $\nu' \in \mathbb{Z}^n$ many times. In general, it is NP-hard to determine whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$ since we must know if $\mathcal{F}_\mathcal{L}(\nu') \neq \emptyset$ and $\mathcal{F}_\mathcal{L}(\nu - \nu') \neq \emptyset$. Instead, we can check a sufficient condition for when $\nu' \notin \mathcal{B}_\mathcal{L}(\nu)$, and so, we compute a superset of a ν -truncated Gröbner basis since we keep some vectors that are not needed.

Firstly, note that in Algorithm 2, whenever we check whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$, we always have $\mathcal{F}_\mathcal{L}(\nu') \neq \emptyset$ since either $\nu' = u^+ \geq 0$ for some $u \in \mathcal{L}$ or $\nu' = \nu^{(u,v)} \geq 0$ for some $u, v \in \mathcal{L}$ and in either case $\nu' \in \mathcal{F}_\mathcal{L}(\nu') \neq \emptyset$. So, in the algorithm, we only need to check whether $\mathcal{F}_\mathcal{L}(\nu - \nu') \neq \emptyset$.

We could instead check for feasibility of a relaxation of the feasible set $\mathcal{F}_\mathcal{L}(\nu - \nu')$. One possible relaxation of $\mathcal{F}_\mathcal{L}(\nu - \nu')$ to check is

$$\mathcal{F}_\mathcal{L}^\mathbb{Z}(\nu - \nu') := \{x : x \equiv \nu - \nu' \pmod{\mathcal{L}}, x \in \mathbb{Z}^n\}.$$

But since $\nu - \nu' \in \mathbb{Z}^n$, we have $\nu - \nu' \in \mathcal{F}_\mathcal{L}^\mathbb{Z}(\nu - \nu') \neq \emptyset$, and so, this is trivially always satisfied. Another possible relaxation is the linear relaxation of $\mathcal{F}_\mathcal{L}(\nu - \nu')$:

$$\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') := \{x : x \equiv \nu - \nu' \pmod{\mathcal{L}^\mathbb{R}}, x \in \mathbb{R}_+^n\}$$

where $\mathcal{L}^\mathbb{R} \subseteq \mathbb{R}^n$ is the smallest subspace containing $\mathcal{L}$; that is, $\mathcal{L}^\mathbb{R} := \{ku : u \in \mathcal{L}, k \in \mathbb{R}\}$. We can thus solve a linear program to check whether $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$ implying that $\mathcal{F}_\mathcal{L}(\nu - \nu') = \emptyset$. Note that $\mathcal{F}_\mathcal{L}(\nu - \nu') = \mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') \cap \mathcal{F}_\mathcal{L}^\mathbb{Z}(\nu - \nu')$, but $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') \neq \emptyset$ and $\mathcal{F}_\mathcal{L}^\mathbb{Z}(\nu - \nu') \neq \emptyset$ do not imply that $\mathcal{F}_\mathcal{L}(\nu - \nu') \neq \emptyset$.

In practice, computational experiments show that it is usually not worthwhile performing the full check whether $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$, so instead, we use a sufficient condition for when $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$ and thus $\nu' \notin \mathcal{B}_\mathcal{L}(\nu)$ that is quick to check. Let

$$\mathcal{L}_+^\circ := \{a \in \mathbb{R}_+^n : au = 0 \ \forall u \in \mathcal{L}\}.$$

Note that $\mathcal{L}_+^\circ$ is a pointed convex cone. Firstly, observe that, for any $a \in \mathcal{L}_+^\circ$ and any $\nu' \in \mathbb{Z}^n$, we have $ax = a\nu'$ for all $x \in \mathcal{F}_\mathcal{L}(\nu')$. Secondly, if $a \in \mathcal{L}_+^\circ$, then, for all $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$, we have $a\nu \geq a\nu'$ since if $x \in \mathcal{F}_\mathcal{L}(\nu - \nu') \neq \emptyset$, then $a(\nu - \nu') = ax$ and $ax \geq 0$ because $a \geq 0$ and $x \geq 0$. Therefore, if $a\nu < a\nu'$, then $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$ and thus $\nu' \notin \mathcal{B}_\mathcal{L}(\nu)$. Moreover, it follows from Farkas' lemma that $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$ if and only if there exists an $a \in \mathcal{L}_+^\circ$ where $a\nu < a\nu'$, and furthermore, $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') = \emptyset$ if and only if there exists an extreme ray a of the cone $\mathcal{L}_+^\circ$ where $a\nu < a\nu'$. The set of extreme rays is finite but there are far too many of them in general to check this condition. So, we need a way of selecting one $a \in \mathcal{L}_+^\circ$ or a small set of a . Choosing different a 's can produce very different results, and the best a 's to choose vary from fiber to fiber.

We now present a novel approach for selecting a *good* $a \in \mathcal{L}_+^\circ$. Now, note that when we run Algorithm 2 and check whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$, we have $a\nu' \geq 0$ for all $a \in \mathcal{L}_+^\circ$ since from above $\mathcal{F}_\mathcal{L}(\nu') \neq \emptyset$ ($\nu' \geq 0$). Ideally, there exists $a \in \mathcal{L}_+^\circ$ where $a\nu = 0$, implying that $a\nu' = 0$ (i.e. if $a_i \neq 0$, then $\nu'_i = 0$) for every $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$. This condition is very strong and is quick to check and effectively means that we compute using a sub-lattice of $\mathcal{L}$. Otherwise if $a\nu > 0$ for all $a \in \mathcal{L}_+^\circ$, then a useful heuristic is to choose a single $a \in \mathcal{L}_+^\circ$ such that $a\nu$ is minimal with respect to some norm $\|\cdot\|$ of a . More formally, we solve the following problem:

$$\operatorname{argmin}\{a\nu : \|a\| = 1, a \in \mathcal{L}_+^\circ\}.$$

If we use the l_1 -norm (i.e. $\|a\|_1 = \sum_i a_i$), then we can find a using linear programming. In this case, note that we only need to solve one linear program to compute a as opposed to solving a linear program every time we check whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$.

Example 4.6. Consider again the set $S := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1)\}$ and the lattice $\mathcal{L} \subseteq \mathbb{Z}^6$ spanned by S . Let $c := (2, 1, 0, 0, 0, 0)$ and let $\prec$ be some term order. Then, a $\prec_c$ -Gröbner basis of $\mathcal{L}$ is

$$G := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1), (1, 1, 5, -1, -3, 0), (0, 1, 3, 1, -2, -1), (2, -1, 1, -5, 0, 3)\}.$$

Recall that this is a $\prec_c$ -Gröbner basis for every possible fiber $\nu \in \mathbb{Z}^n$. We now examine truncated $\prec_c$ -Gröbner basis for two different fibers.

- (i). Consider $\nu := (0, 1, 0, 5, 0, 0)$ (see Figure 2b); $\mathcal{F}_\mathcal{L}(\nu) = \{(2, 0, 1, 0, 0, 3), (0, 1, 0, 5, 0, 0)\}$. Since the feasible set consists of only two feasible solutions, the minimal ν -truncated $\prec_c$ -Gröbner basis contains only one vector: the vector between the two feasible solutions. Thus, the set $\{(2, -1, 1, -5, 0, 3)\}$ is a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$.

If we run the truncated completion procedure, Algorithm 2, using $\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu') \neq \emptyset$ as a check for truncation, we compute the set $\mathcal{CP}_\nu(\succ, S) = \{(2, -1, 1, -5, 0, 3)\}$.

If instead we run the truncated completion procedure using the quick truncation test $a\nu < a\nu'$ where $a := \frac{1}{4}(0, 1, 1, 0, 2, 0) = \operatorname{argmin}\{a\nu : \|a\|_1 = 1, a \in \mathcal{L}_+^\circ\}$, we again compute the set $\mathcal{CP}_\nu(\succ, S) = \{(2, -1, 1, -5, 0, 3)\}$. Note that $a\nu = \frac{1}{4}(0, 1, 1, 0, 2, 0) \cdot (0, 1, 0, 5, 0, 0) = \frac{1}{4}$. Then, for example, $a\nu' = \frac{1}{4}(0, 1, 1, 0, 2, 0) \cdot (1, -1, -1, -3, 1, 2)^+ = \frac{1}{2} > a\nu$; hence, the vector $(1, -1, -1, -3, 1, 2)$ is not needed in a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$.

On the other hand, if we used the vector $a := \frac{1}{14}(0, 0, 1, 5, 0, 8) \in \mathcal{L}_+^\circ$, then the quick truncation check $a\nu < a\nu'$ is useless, and we would compute all five vectors of the $\prec_c$ -Gröbner basis of $\mathcal{L}$.

- (ii). Consider $\nu := (2, 2, 4, 2, 0, 0)$ (see Figure 2a); $\mathcal{F}_{\mathcal{L}}(\nu) = \{(2, 1, 1, 1, 2, 1), (2, 2, 4, 2, 0, 0)\}$. Since, the feasible set consists of only two feasible solutions, the set $\{(0, 1, 3, 1, -2, -1)\}$ is a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$.

Using $\mathcal{F}_{\mathcal{L}}^{\mathbb{R}}(\nu - \nu') \neq \emptyset$ as a check for truncation, we compute the following ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$: $G := \{(0, 1, 3, 1, -2, -1), (1, 0, 2, -2, -1, 1)\}$. So, we have computed an additional vector $(1, 0, 2, -2, -1, 1)$ that is not strictly needed since $\mathcal{F}_{\mathcal{L}}(\nu - \nu') = \emptyset$ even though $\mathcal{F}_{\mathcal{L}}^{\mathbb{R}}(\nu - \nu') \neq \emptyset$ where $\nu' := (1, 0, 2, -2, -1, 1)^+ = (1, 0, 2, 0, 0, 1)$.

Using the vector $a := \frac{1}{4}(0, 1, 0, 1, 0, 2) = \operatorname{argmin}\{a\nu : \|a\|_1 = 1, a \in \mathcal{L}_+^{\circ}\}$ as a quick check for truncation, we obtain the following ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$:

$$G := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1), (1, 1, 5, -1, -3, 0), (0, 1, 3, 1, -2, -1)\}.$$

So here, we have computed three additional unnecessary vectors. If instead we use the vector $\frac{1}{14}(0, 0, 1, 5, 0, 8)$, then we would have computed only three vectors. In this case, there is no single vector in $a \in \mathcal{L}_+^{\circ}$ that results in only two vectors being computed.

The following example demonstrates the potential speed increase from computing a truncated Gröbner basis as opposed to computing a full Gröbner basis. We use three different methods for checking whether $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$ in order of increasing effectiveness:

- (i). $a\nu < a\nu'$ where $a = \operatorname{argmin}\{a\nu : \|a\|_1 = 1, a \in \mathcal{L}_+^{\circ}\}$,
- (ii). $\mathcal{F}_{\mathcal{L}}^{\mathbb{R}}(\nu - \nu') = \emptyset$, or
- (iii). $\mathcal{F}_{\mathcal{L}}(\nu - \nu') = \emptyset$.

Since criterion (i) is in general much faster to check than (ii), we always check (i) before (ii), and similarly, since criterion (i) and (ii) are in general much faster to check than (iii), we always check them both before applying criterion (iii). We solve $\mathcal{F}_{\mathcal{L}}^{\mathbb{R}}(\nu - \nu') = \emptyset$ using the simplex algorithm implementation in the GLPK (GNU Linear Programming Kit) package.¹ We solve $\mathcal{F}_{\mathcal{L}}(\nu - \nu') = \emptyset$ using the branch-and-bound implementation in GLPK, which is not useful in practice, but we use it here to show the sizes of minimal truncated Markov bases.

Example 4.7. Let $\mathcal{L} = \mathcal{L}_A := \{u : Au = \mathbf{0}, u \in \mathbb{Z}^n\}$ where

$$A = \begin{bmatrix} 15 & 4 & 14 & 19 & 2 & 1 & 10 & 17 & 11 & 9 & 4 & 15 & 20 \\ 18 & 11 & 13 & 5 & 16 & 16 & 8 & 19 & 18 & 21 & 5 & 7 & 1 \\ 11 & 7 & 8 & 19 & 15 & 18 & 14 & 6 & 1 & 23 & 11 & 3 & 10 \\ 17 & 10 & 13 & 17 & 16 & 14 & 15 & 18 & 3 & 2 & 1 & 17 & 1 \end{bmatrix}$$

The size of a minimal Markov basis of $\mathcal{L}$ is 10868. Let $c = (3, 15, 1, 5, 2, 17, 16, 16, 15, 9, 7, 11, 13)$. The size of a minimal $\prec_c$ -Gröbner basis of $\mathcal{L}$ for some term order $\prec$ is 24941. This takes 106.96 seconds to compute using 4ti2. In the following table, we list the time taken to compute truncated Gröbner bases from the minimal Markov basis of $\mathcal{L}$. The first column lists the values used for ν . In the following columns, we list the size of the computed set and the time taken for each of the three possible ways to check whether $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$.

¹GLPK is open source and freely available from <http://www.gnu.org/software/glpk/>

ν	$a \in \mathcal{L}_+^\circ$		$\mathcal{F}_{\mathcal{L}}^{\text{rk}}(\nu - \nu'')$		$\mathcal{F}_{\mathcal{L}}(\nu - \nu'')$	
(1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 0)	307	0.02s	1	0.09s	0	0.10s
(1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)	418	0.10s	36	0.19s	0	0.31s
(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	6494	1.61s	201	1.20s	0	2.18s
(1, 2, 0, 3, 5, 0, 1, 3, 0, 4, 0, 1, 0)	12191	6.31s	5028	4.43s	158	186.81s
(19, 7, 3, 8, 13, 11, 1, 15, 4, 8, 17, 9, 5)	24748	108.19s	24334	107.25s	24284	> 3600s

For the example above, choosing a single $a \in \mathcal{L}_+^\circ$ where $a := \operatorname{argmin}\{a\nu : \|a\|_1 = 1, a \in \mathcal{L}_+^\circ\}$ works reasonably well when used for checking for truncation. However, in general, using more than one such a may be significantly better particularly when the support of a (the set of non-zero components) is small.

Observe that to compute a truncated Gröbner bases in the previous example, we first needed to compute a Markov basis, and in some cases, computing the Markov basis took significantly longer than computing the truncated Gröbner basis. This provides motivation for the next section in which we compute truncated Markov basis.

5 Computing truncated Markov bases

In this section, we give a Project-and-Lift algorithm for computing truncated Markov bases.

Given $\sigma \subseteq \{1, \dots, n\}$, we define the projective map $\pi_\sigma : \mathbb{Z}^n \mapsto \mathbb{Z}^{|\sigma|}$ that projects a vector in $\mathbb{Z}^n$ onto the σ components. We define $\mathcal{L}^\sigma$ where $\sigma \subseteq \{1, \dots, n\}$ as the projection of $\mathcal{L}$ onto the σ components – that is, $\mathcal{L}^\sigma = \pi_\sigma(\mathcal{L})$. Note that $\mathcal{L}^\sigma$ is also a lattice. For ease of notation, we will often denote the singleton set $\{i\}$ as just i , and so, for example, $\mathcal{L}^{\{i\}}$ is denoted $\mathcal{L}^i$ and $\pi_{\{i\}}$ is denoted π_i . It should be clear from the context whether by i we mean $\{i\}$ or just i .

The fundamental idea behind the Project-and-Lift algorithm is that, for some $\nu \in \mathbb{Z}^n$, using a set $S \subseteq \mathcal{L}^i$ that is a $\pi_i(\nu)$ -truncated Markov basis of $\mathcal{L}^i$ for some $i \in \{1, \dots, n\}$, we can compute a set $S' \subseteq \mathcal{L}^i$ such that S' lifts to a ν -truncated Markov basis of $\mathcal{L}$. So, for some $\sigma \subseteq \{1, \dots, n\}$, since $\mathcal{L}^\sigma$ is also a lattice, starting with a ν_σ -truncated Markov basis of $\mathcal{L}^\sigma$, we can compute a $\pi_{\sigma \setminus i}(\nu)$ -truncated Markov basis of $\mathcal{L}^{\sigma \setminus i}$ for some $i \in \sigma$. By doing this repeatedly for every $i \in \sigma$, we attain a ν -truncated Markov basis of $\mathcal{L}$.

First, we extend the definition of reduction paths and ν -truncated Gröbner bases. Given some vector $c \in \mathbb{Q}^n$, a path $(x^0, \dots, x^k)$ in $\mathcal{G}_{\mathcal{L}}(\nu, G)$ is an c -reduction path if for all $j \in \{1, \dots, k-1\}$, we have either $cx^0 \geq cx^j$ or $cx^k \geq cx^j$. A set $G \subseteq \mathcal{L}$ is a **c -Gröbner basis** of $\mathcal{F}_{\mathcal{L}}(\nu)$ if for every pair $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$, there exists a c -reduction path from x to y in $\mathcal{G}_{\mathcal{L}}(\nu, G)$. A set $G \subseteq \mathcal{L}$ is a ν -truncated **c -Gröbner basis** of $\mathcal{L}$ if for all $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$, G is a c -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu')$.

The following lemma is fundamental to the Project-and-Lift algorithm. Note that the property that $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$ for some $i \in \{1, \dots, n\}$ means that the map π_i from $\mathcal{L}$ to $\mathcal{L}^i$ is a bijection and thus, the inverse map $\pi_i^{-1} : \mathcal{L}^i \mapsto \mathcal{L}$ is well-defined (each vector in $\mathcal{L}^i$ lifts to a unique vector in $\mathcal{L}$). Moreover, by linear algebra, for all $u \in \mathcal{L}$, there must exist a vector $\omega^i \in \mathbb{Q}^{n-1}$ such that $\omega^i \cdot \pi_i(u) = u_i$. We always write such a vector as ω^i . Importantly, note that given $\nu \in \mathbb{Z}^n$, we have $\pi_i(\mathcal{B}_{\mathcal{L}}(\nu)) \subseteq \mathcal{B}_{\mathcal{L}^i}(\pi_i(\nu))$ since if $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$, then there exists $\gamma \in \mathcal{F}_{\mathcal{L}}(\nu - \nu')$, and so, $\pi_i(\gamma) \in \mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu - \nu')) \neq \emptyset$, and consequently, $\pi_i(\nu') \in \mathcal{B}_{\mathcal{L}^i}(\pi_i(\nu))$.

Lemma 5.1. *Let $i \in \{1, \dots, n\}$ where $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$, and let $S \subseteq \mathcal{L}^i$. Let $\nu \in \mathbb{Z}^n$. If S is a $\pi_i(\nu)$ -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}^i$, then $\pi_i^{-1}(S)$ is a ν -truncated $(-e^i)$ -Gröbner basis of $\mathcal{L}$.*

Proof. Assume S is a $\pi_i(\nu)$ -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}^i$. Let $x, y \in \mathcal{F}_{\mathcal{L}}(\nu')$ for some $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$. We need to show that there is an $(-e^i)$ -reduction path from x to y in $\mathcal{G}_{\mathcal{L}}(\nu', \pi_i^{-1}(S))$. Let $\tilde{x} = \pi_i(x)$, $\tilde{y} = \pi_i(y)$, and $\tilde{\nu}' = \pi_i(\nu') \in \mathcal{B}_{\mathcal{L}^i}(\pi_i(\nu))$. By assumption, there exists an $(-\omega^i)$ -reduction path $(\tilde{x} = \tilde{x}^0, \dots, \tilde{x}^k = \tilde{y})$ in $\mathcal{G}_{\mathcal{L}^i}(\tilde{\nu}', S)$. So, we have either $\omega^i \tilde{x}^j \geq \omega^i \tilde{x}$ or $\omega^i \tilde{x}^j \geq \omega^i \tilde{y}$ for all j . We now lift this $(-\omega^i)$ -reduction path in $\mathcal{G}_{\mathcal{L}^i}(\tilde{\nu}', S)$ to an $(-e^i)$ -reduction path in $\mathcal{G}_{\mathcal{L}}(\nu', \pi_i^{-1}(S))$. Let $x^j = \nu' + \pi_i^{-1}(\tilde{x}^j - \tilde{\nu}')$ for all $j = 0, \dots, k$. Hence, $\pi_i(x^j) = \tilde{x}^j$ and $\omega^i x^j = x_i^j$, and therefore, either $x_i^j \geq x_i$ or $x_i^j \geq y_i$. Also, $x^j - x^{j-1} = \pi_i^{-1}(\tilde{x}^j - \tilde{x}^{j-1}) \in \pi_i^{-1}(S)$ for all $j = 1, \dots, k$. Therefore, $(x = x^0, \dots, y^k = y)$ is an $(-e^i)$ -reduction path in $\mathcal{G}_{\mathcal{L}}(\nu, \pi_i^{-1}(S))$ as required. $\square$

The converse of Lemma 5.1 is not true: if $\pi_i^{-1}(S)$ is a ν -truncated $(-e^i)$ -Gröbner basis of $\mathcal{L}$, then S is *not necessarily* a $\pi_i(\nu)$ -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}$. If $\pi_{\sigma}(\mathcal{B}_{\mathcal{L}}(\nu)) = \mathcal{B}_{\mathcal{L}^{\sigma}}(\nu_{\sigma})$, then the converse holds, since if $\pi_i^{-1}(S)$ is a $(-e^i)$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\nu')$ for some $\nu' \in \mathbb{Z}^n$, then S is a $(-\omega^i)$ -Gröbner basis of $\mathcal{F}_{\mathcal{L}}(\bar{\nu}')$ where $\bar{\nu}' = \pi_i(\nu')$ because $(-e^i)$ -reduction paths in $\mathcal{G}_{\mathcal{L}}(\nu', \pi_i^{-1}(S))$ project to $(-\omega^i)$ -reduction paths in $\mathcal{G}_{\mathcal{L}^i}(\bar{\nu}', S)$. In general however, $\pi_i(\mathcal{B}_{\mathcal{L}}(\nu)) \subseteq \mathcal{B}_{\mathcal{L}^i}(\pi_i(\nu))$, and we may have $\pi_i(\mathcal{B}_{\mathcal{L}}(\nu)) \subsetneq \mathcal{B}_{\mathcal{L}^i}(\pi_i(\nu))$.

By definition, an $(-e^i)$ -Gröbner basis of $\mathcal{L}$ is a Markov basis of $\mathcal{L}$. Conversely, a Markov basis of $\mathcal{L}$ is also an $(-e^i)$ -Gröbner basis of $\mathcal{L}$. This follows since, given a Markov basis of $\mathcal{L}$, for any $x, y \in \mathcal{F}_{\mathcal{L}}(\nu)$ for any ν , there must exist a path from $x - \gamma$ to $y - \gamma$ where $\gamma = \min\{x_i, y_i\} \cdot e^i$, and by translating such a path by γ , we get an $(-e^i)$ -reduction path from x to y . So, we arrive at the following corollary.

Corollary 5.2. *Let $i \in \{1, \dots, n\}$ where $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$, and let $S \subseteq \mathcal{L}^i$. If S is a $\pi_i(\nu)$ -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}^i$, then $\pi_i^{-1}(S)$ is a ν -truncated Markov basis of $\mathcal{L}$.*

Given any vector $c \in \mathbb{Z}^n$ and a term order $\prec$ for $\mathcal{L}$, recall that for the ordering $\prec_c$, we have $x \prec_c y$ if $cx < cy$ or $cx = cy$ and $x \prec y$. Also, recall that the order $\prec_c$ is a term order if and only if $IP_{\mathcal{L}, c}(\mathbf{0}) = 0$. Importantly then, a $\prec_c$ -reduction path is also a c -reduction path. So, we can compute a ν -truncated c -Gröbner basis by computing a ν -truncated $\prec_c$ -Gröbner basis.

If $IP_{\mathcal{L}, -e^i}(\mathbf{0}) := \max\{x_i : x \in \mathcal{L}, x \in \mathbb{N}^n\} = 0$, we say that i is **bounded** for $\mathcal{L}$ and **unbounded** otherwise. Thus, $\prec_{-e^i}$ is a term order for $\mathcal{L}$ if and only if i is bounded, and moreover, $\prec_{-\omega^i}$ is a term order for $\mathcal{L}^i$ if and only if i is bounded since

$$IP_{\mathcal{L}, -e^i}(\mathbf{0}) = \max\{x_i : x \in \mathcal{L}, x \in \mathbb{N}^n\} = \max\{\omega^i x : x \in \mathcal{L}^i, x \in \mathbb{N}^{n-1}\} = IP_{\mathcal{L}^i, -\omega^i}(\mathbf{0}).$$

Now if i is bounded, then the ordering $\prec_{-\omega^i}$ is a term order for $\mathcal{L}^i$, and so, given a set $S \subseteq \mathcal{L}^i$ that is a ν -truncated Markov basis of $\mathcal{L}^i$, we can compute a ν -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}^i$ using Algorithm 2. In other words, the set $S' = \mathcal{CP}_{\pi_i(\nu)}(\prec_{-\omega^i}, S)$ is a $\pi_i(\nu)$ -truncated $(-\omega^i)$ -Gröbner basis of $\mathcal{L}^i$, and by Corollary 5.2, the set $\pi_i^{-1}(S')$ is a ν -truncated Markov basis of $\mathcal{L}$.

If i is unbounded, then computing a ν -truncated Markov basis of $\mathcal{L}$ from a $\pi_i(\nu)$ -truncated Markov basis of $\mathcal{L}^i$ is actually more straight-forward than otherwise. Crucially, i is unbounded if and only if $IP_{\mathcal{L}, -e^i}(\mathbf{0}) > 0$ or equivalently there exists $u \in \mathcal{L} \cap \mathbb{N}^n$ where $u_i > 0$. Then assuming $\ker(\pi_i) \cap \mathcal{L} = \emptyset$, given a set $S \subseteq \mathcal{L}^i$ that is a $\pi_i(\nu)$ -truncated Markov basis of $\mathcal{L}^i$, it suffices to add u to $\pi_i^{-1}(S')$ to create a ν -truncated Markov basis of $\mathcal{L}$ (see Lemma 5.3 below). We can use linear programming to check whether i is unbounded and also to find such a $u \in \mathcal{L} \cap \mathbb{N}^n$ where $u_i > 0$.

Lemma 5.3. *Let $i \in \{1, \dots, n\}$ and $\ker(\pi_i) \cap \mathcal{L} = \emptyset$, and $u \in \mathcal{L} \cap \mathbb{N}^n$ where $u_i > 0$. If $S \subseteq \mathcal{L}^i$ is a $\pi_i(\nu)$ -truncated Markov basis of $\mathcal{L}^i$, then $\pi_i^{-1}(S) \cup \{u\}$ is a ν -truncated Markov basis of $\mathcal{L}$.*

Proof. Let $x', y' \in \mathcal{F}_{\mathcal{L}}(\nu')$ for some $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$. Since S is a $\pi_i(\nu)$ -truncated Markov basis of $\mathcal{L}^i$, there exists a path from $\pi_i(x')$ to $\pi_i(y')$. We can convert this path into a $(-\omega^i)$ -reduction path in $\mathcal{G}_{\mathcal{L}^i}(\pi_i(\nu'), S \cup \pi_i(u))$ by adding $\pi_i(u)$ to the start of the path as many times as necessary and subtracting $\pi_i(u)$ from the end of the path the same number of times. This works since $\omega^i(\pi_i(u)) = u_i > 0$ and $u \geq 0$. As in Lemma 5.1, we can then lift this to a path from x' to y' in $\mathcal{G}_{\mathcal{L}}(\nu', \pi_i^{-1}(S) \cup \{u\})$. $\square$

We can apply the above reasoning to compute a Markov basis of $\mathcal{L}^{\sigma \setminus i}$ from a Markov basis of $\mathcal{L}^{\sigma}$ for some $\sigma \subseteq \{1, \dots, n\}$ and $i \in \sigma$. First, analogously to π_i and $\prec_{-\omega^i}$ in the context of $\mathcal{L}^i$ and $\mathcal{L}$, we define π_i^{σ} and $\prec_{-\omega^i}^{\sigma}$ in the same way except in the context of $\mathcal{L}^{\sigma}$ and $\mathcal{L}^{\sigma \setminus i}$ respectively.

We can now present our Project-and-Lift algorithm (Algorithm 3).

Algorithm 3 Project-and-Lift algorithm

Input: a lattice $\mathcal{L}$ and a vector $\nu \in \mathbb{Z}^n$.

Output: a ν -truncated Markov basis M of $\mathcal{L}$

Find a set $\sigma \subseteq \{1, \dots, n\}$ such that $\ker(\pi_{\sigma}) \cap \mathcal{L} = \{\mathbf{0}\}$.

Compute a set $M \subseteq \mathcal{L}^{\sigma}$ that is a ν_{σ} -truncated Markov basis of $\mathcal{L}^{\sigma}$.

while $\sigma \neq \emptyset$ **do**

 Select $i \in \sigma$

if i is bounded **then**

$G := \mathcal{CP}_{\nu_{\sigma}}(\prec_{-\omega^i}^{\sigma}, M)$

$M := (\pi_i^{\sigma})^{-1}(G)$

else

 Compute $u \in \mathcal{L}^{\sigma \setminus i} \cap \mathbb{N}^n$ such that $u_i > 0$

$M := (\pi_i^{\sigma})^{-1}(M) \cup \{u\}$

end if

$\sigma := \sigma \setminus i$

$M := \{u \in M : u^+ \in \mathcal{B}_{\mathcal{L}^{\sigma}}(\nu_{\sigma})\}$

end while

return M .

Lemma 5.4. *Algorithm 3 terminates and satisfies its specifications.*

Proof. Algorithm 3 terminates, since Algorithm 2, which computes $\mathcal{CP}_{\nu_{\sigma}}(\prec_{-\omega^i}^{\sigma}, M)$, always terminates.

We claim that for each iteration of the algorithm, M is a ν_{σ} -truncated Markov basis of $\mathcal{L}^{\sigma}$ and $\ker(\pi_{\sigma}) \cap \mathcal{L} = \{\mathbf{0}\}$; therefore, at termination, M is a ν -truncated Markov basis of $\mathcal{L}$. This is true for the first iteration, so we assume it is true for the current iteration.

If $\sigma = \emptyset$, then there is nothing left to do, so assume otherwise. Since by assumption, $\ker(\pi_{\sigma}) \cap \mathcal{L}^{\sigma} = \{\mathbf{0}\}$, we must have $\ker(\pi_i^{\sigma}) \cap \mathcal{L}^{\sigma \setminus i} = \{\mathbf{0}\}$, and so, the inverse map $(\pi_i^{\sigma})^{-1} : \mathcal{L}^{\sigma} \rightarrow \mathcal{L}^{\sigma \setminus i}$ is well-defined. Let $i \in \sigma$, and $\sigma' := \sigma \setminus i$. If i is bounded, then let $G := \mathcal{CP}_{\nu_{\sigma}}(\prec_{-\omega^i}^{\sigma}, M)$; then, G is a $\prec_{-\omega^i}^{\sigma}$ -Gröbner basis of $\mathcal{L}^{\sigma}$. Let $M' := (\pi_i^{\sigma})^{-1}(G)$, and then by Corollary 5.2, M' is a $\pi_{\sigma'}(\nu)$ -truncated Markov basis of $\mathcal{L}^{\sigma'}$. Otherwise, let $M := (\pi_i^{\sigma})^{-1}(M) \cup \{u\}$ where $u \in \mathcal{L}^{\sigma \setminus i} \cap \mathbb{N}^n$ such that $u_i > 0$, and by Lemma 5.3, M' is a $\pi_{\sigma'}(\nu)$ -truncated Markov basis of $\mathcal{L}^{\sigma'}$. Also, $M'' := \{u \in M' : u^+ \in \mathcal{B}_{\mathcal{L}^{\sigma'}}(\pi_{\sigma'}(\nu))\}$ must be a $\pi_{\sigma'}(\nu)$ -truncated Markov basis of $\mathcal{L}^{\sigma'}$.

Lastly, since $\sigma' \subseteq \sigma$, we must have $\ker(\pi_{\sigma'}) \cap \mathcal{L} = \{\mathbf{0}\}$. Thus, the claim is true for the next iteration. $\square$

Initially in our Project-and-Lift algorithm, we need to find a set $\sigma \subseteq \{1, \dots, n\}$ such that $\ker(\pi_\sigma) \cap \mathcal{L}^\sigma = \{\mathbf{0}\}$, and then, we need to compute a Markov basis for $\mathcal{L}^\sigma$. This is actually quite straightforward and can be done in polynomial time. Let B be a basis for the lattice $\mathcal{L}$ ($\mathcal{L}$ is spanned by the rows of the matrix B). Let $k := \text{rank}(B)$. Any k linearly independent columns of B then suffice to give a set $\bar{\sigma}$ such that every vector in $\mathcal{L}^\sigma$ lifts to a unique vector in $\mathcal{L}$; that is, $\ker(\pi_{\bar{\sigma}}) \cap \mathcal{L}^\sigma = \{\mathbf{0}\}$. Such a set $\bar{\sigma}$ can be found via Gaussian elimination. Let $S = \pi_{\bar{\sigma}}(B)$; then, S spans $\mathcal{L}^\sigma$, and $S \in \mathbb{Z}^{k \times k}$ since $|\bar{\sigma}| = k$. Let S' be an upper triangle matrix with positive diagonal entries and non-positive entries elsewhere such that (the rows of) S' span $\mathcal{L}^\sigma$. We can always construct such a matrix S' from S in polynomial time using the *Hermite Normal Form* (HNF) algorithm (see for example [14]). Also, S' is a Markov basis of $\mathcal{L}^\sigma$ since it is actually a Gröbner basis of $\mathcal{L}^\sigma$ with respect to a *lexicographic* ordering, and thus, a Markov basis of $\mathcal{L}^\sigma$.

Example 5.5. Consider again the set $S := \{(1, -1, -1, -3, 1, 2), (1, 0, 2, -2, -1, 1)\}$, and the lattice $\mathcal{L} \subseteq \mathbb{Z}^6$ spanned by S . Let again $\nu := (0, 1, 0, 5, 0, 0)$ (see Figure 2b). Recall that the minimal ν -truncated $\prec_c$ -Gröbner basis is $\{(2, -1, 1, -5, 0, 3)\}$, since $\mathcal{F}_{\mathcal{L}}(\nu) = \{(2, 0, 1, 0, 0, 3), (0, 1, 0, 5, 0, 0)\}$.

Let $\sigma = \{3, 4, 5, 6\}$. Then, $\ker(\pi_\sigma) \cap \mathcal{L}^\sigma = \{\mathbf{0}\}$. Note that $\pi_\sigma(S) = \{(1, -1), (1, 0)\}$. The set $M = \{(1, 0), (0, 1)\}$ is a Markov basis of $\mathcal{L}^\sigma$.

- (i). Set $i := 3$. Then, i is unbounded since $(1, 0, 2) \in \mathcal{L}^{\sigma \setminus i}$. Thus, $M = \{(1, 0, 2), (0, 1, 3)\}$ is a Markov basis of $\mathcal{L}^{\sigma \setminus i}$. Set $\sigma = \{4, 5, 6\}$.
- (ii). Set $i := 5$. Then, i is bounded. $M = \{(-2, 1, -1), (1, 0, 2)\}$ is a minimal $(0, 1, 0)$ -truncated $\prec_{-\omega^i}$ -Gröbner basis of $\mathcal{L}^\sigma$. Note that $\nu_{\bar{\sigma}} = (0, 1, 0)$. So, $M = \{(-2, 1, -1, 0), (1, 0, 2, -1)\}$ is a $(0, 1, 0, 0)$ -truncated Markov basis of $\mathcal{L}^{\sigma \setminus i}$. Set $\sigma = \{4, 6\}$.
- (iii). Set $i := 4$. Then, i is bounded. $M = \{(2, -1, 1, 0)\}$ is a minimal $(0, 1, 0, 0)$ -truncated $\prec_{-\omega^i}$ -Gröbner basis of $\mathcal{L}^\sigma$. So, $M = \{(2, -1, 1, -5, 0)\}$ is a $(0, 1, 0, 5, 0)$ -truncated Markov basis of $\mathcal{L}^{\sigma \setminus i}$. Set $\sigma = \{6\}$.
- (iv). Set $i := 6$. Then, i is bounded. $M = \{(-2, 1, -1, 5, 0)\}$ is a minimal $(0, 1, 0, 5, 0)$ -truncated $\prec_{-\omega^i}$ -Gröbner basis of $\mathcal{L}^\sigma$. So, $M = \{(-2, 1, -1, 5, 0, -3)\}$ is a $(0, 1, 0, 5, 0, 0)$ -truncated Markov basis of $\mathcal{L}$.

In the above Markov basis computation, the size of the set M was never larger than 2 although the size of a full minimal Markov basis is 5.

In the next example, we show the computational benefits of computing a truncated Markov basis as opposed to the full Markov basis.

Example 5.6. Let $\mathcal{L} = \mathcal{L}_A$ for the matrix A given in Example 4.7. The size of a minimal Markov basis of $\mathcal{L}$ is 10868. It takes 36.39 seconds to compute.

Let $\nu = (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$. The size of a minimal ν -truncated Markov basis of $\mathcal{L}$ is 0. In the next table, we list the times taken to compute a ν -truncated Markov basis using the three different criteria for truncation. We also list the sizes of the intermediate Gröbner basis computations. The number at the top of the column is the size of σ as used in algorithm 3. The first row is the case without using truncation. Note how the intermediate sizes of the truncated computations remain much smaller than the final size of a full Markov basis of $\mathcal{L}$.

Truncation	4	3	2	1	0	Time
none	545	1822	3681	12573	10868	36.39s
$a \in \mathcal{L}_+^\circ$	545	977	1302	1846	564	1.20s
$\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu'')$	545	977	878	697	194	2.16s
$\mathcal{F}_\mathcal{L}(\nu - \nu'')$	545	6	3	4	0	20.73s

In the following table, we list the times taken to compute a minimal truncated Markov basis for different ν . The first column lists the values used for ν . In the next columns, we list the size of the computed set and the time taken for each of the three possible ways to check whether $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$.

ν	$a \in \mathcal{L}_+^\circ$		$\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu'')$		$\mathcal{F}_\mathcal{L}(\nu - \nu'')$	
(1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0)	4	0.76s	1	0.90s	0	3.33s
(1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)	158	16.70s	36	21.28s	0	> 3600s
(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	546	1.20s	194	2.16s	0	20.73s
(1, 2, 0, 3, 5, 0, 1, 3, 0, 4, 0, 1, 0)	7381	7.61s	3734	11.47s	146	> 3600s
(19, 7, 3, 8, 13, 11, 1, 15, 4, 8, 17, 9, 5)	10814	39.04s	10761	44.07s	10739	> 3600s

Not only have we reduced the time to compute a Markov basis, we also have reduced the time to compute a truncated Gröbner basis by using a truncated Markov basis instead of a full Markov basis. The following table lists the times for computing a truncated Gröbner basis from a truncated Markov basis using the same cost vector $c = (3, 15, 1, 5, 2, 17, 16, 16, 15, 9, 7, 11, 13)$ as before.

ν	$a \in \mathcal{L}_+^\circ$		$\mathcal{F}_\mathcal{L}^\mathbb{R}(\nu - \nu'')$		$\mathcal{F}_\mathcal{L}(\nu - \nu'')$	
(1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 0)	4	0.00s	1	0.00s	0	0.00s
(1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)	167	0.00s	36	0.19s	0	0.00s
(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	844	0.05s	201	0.11s	0	0.00s
(1, 2, 0, 3, 5, 0, 1, 3, 0, 4, 0, 1, 0)	11768	5.92s	5028	4.07s	158	5.98s
(19, 7, 3, 8, 13, 11, 1, 15, 4, 8, 17, 9, 5)	24729	109.57	24334	107.18s	24284	> 3600s

Observe that in the previous example, the size of a truncated Markov basis for the fiber where $\nu = (1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)$ was very small, but it took much longer to compute than a much larger truncated Markov basis of other fibers. This anomaly can be attributed to the order in which the variables are chosen during the project-and-lift algorithm. If we reorder the variables so that the zero components in $\nu = (1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)$ are chosen first, then the algorithm computes a truncated Markov basis much faster. Hence, the efficiency of the algorithm is sensitive to the order in which the variables are chosen; therefore, future experimentation is needed to determine a heuristic for choosing a *good* variable ordering.

6 Feasibility

The project-and-lift algorithm can be used to find a feasible solution of $\mathcal{F}_\mathcal{L}(\nu)$ for any $\nu \in \mathbb{Z}^n$. Alternatively, in the truncated case, given a particular ν , the project-and-lift algorithm can be used to find a feasible solution of $\mathcal{F}_\mathcal{L}(\nu)$ and also a feasible solution of $\mathcal{F}_\mathcal{L}(\nu')$ for any $\nu' \in \mathcal{B}_\mathcal{L}(\nu)$. Thus, at the same time as finding a Markov basis, the algorithm can also compute a feasible solution.

The basic idea is that, given some $\nu \in \mathbb{Z}^n$ and a feasible solution of $\mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$ for some $i \in \{1, \dots, n\}$ where $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$, we can construct a feasible solution of $\mathcal{F}_\mathcal{L}(\nu)$ if such a feasible

solution exists. Hence, for some $\sigma \subseteq \{1, \dots, n\}$, starting with a feasible solution of $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$ (we choose σ such that this feasible solution is easy to find), we can compute a feasible solution of $\mathcal{F}_{\mathcal{L}^{\sigma \setminus i}}(\pi_{\sigma \setminus i}(\nu))$ for some $i \in \sigma$. By doing this repeatedly for every $i \in \sigma$, we attain a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$.

We now show how to construct a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ from a feasible solution of $\mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$ for some $i \in \{1, \dots, n\}$ where $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$. Let $\nu \in \mathbb{Z}^n$, and $x \in \mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$. First, we lift x . Let $x' := \pi_i^{-1}(x - \pi_i(\nu)) + \nu$. Note that $(x - \pi_i(\nu)) \in \mathcal{L}^i$ and so π_i^{-1} is well-defined. Now, we have $\pi_i(x') = x$ and so $x'_j \geq 0$ for all $j \neq i$. Also, $x' \in \mathcal{F}_{\mathcal{L}}^{\mathbb{Z}}(\nu)$. If $x'_i \geq 0$, then $x' \in \mathcal{F}_{\mathcal{L}}(\nu)$ and we are done. So, assume $x'_i < 0$. Now, if i is unbounded, then there exists $u \in \mathcal{L} \cap \mathbb{N}^n$ where $u_i > 0$; therefore, $x' + \lambda u$ for some $\lambda \in \mathbb{N}$ is non-negative on the i th component and thus a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$. If i is bounded, then we can compute a set G that is a $\pi_i(\nu)$ -truncated $\prec_{-\omega^i}$ -Gröbner basis of $\mathcal{L}^i$ for some term order $\prec$. Let $x'' = \mathcal{NF}(x, G)$. Hence, x'' is the optimal solution of $IP_{\mathcal{L}^i, \prec_{-\omega^i}}(\pi_i(\nu))$ and so also an optimal solution of $IP_{\mathcal{L}^i, -\omega^i}(\pi_i(\nu))$. Thus conceptually, when computing $\mathcal{NF}(x, G)$, we are just maximising the i th component which will thus become non-negative if a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ exists. Therefore, from Lemma 6.1 below, either $x' := \pi_i^{-1}(x'' - \pi_i(\nu)) + \nu$ is a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ or $\mathcal{F}_{\mathcal{L}}(\nu) = \emptyset$.

Lemma 6.1. *Let $i \in \{1, \dots, n\}$ where i is bounded and $\ker(\pi_i) \cap \mathcal{L} = \{\mathbf{0}\}$. Let $\nu \in \mathbb{Z}^n$, and let $x \in \mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$ be a $(-\omega^i)$ -minimal solution of $\mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$. Then, $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$ if and only if $(\pi_i^{-1}(x - \pi_i(\nu)) + \nu) \in \mathcal{F}_{\mathcal{L}}(\nu)$.*

Proof. Let $x' = (\pi_i^{-1}(x - \pi_i(\nu)) + \nu)$. If $x' \in \mathcal{F}_{\mathcal{L}}(\nu)$, then $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$ by definition. We now prove the converse. Assume $\mathcal{F}_{\mathcal{L}}(\nu) \neq \emptyset$, and let $x'' \in \mathcal{F}_{\mathcal{L}}(\nu)$. Then, $\pi_i(x'') \in \mathcal{F}_{\mathcal{L}^i}(\pi_i(\nu))$ and so $-\omega^i \pi_i(x'') \geq -\omega^i x$ which implies that $x''_i \leq (\pi_i^{-1}(x''))_i$. Therefore, $\pi_i^{-1}(x'')$ is non-negative and a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$. $\square$

To compute a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$, we need to start from a feasible solution of $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$ for some $\sigma \subseteq \{1, \dots, n\}$ where $\ker(\pi_\sigma) \cap \mathcal{L} = \{\mathbf{0}\}$. As before in Section 5, we can find a set $\sigma \subseteq \{1, \dots, n\}$ such that $\ker(\pi_\sigma) \cap \mathcal{L}^\sigma = \{\mathbf{0}\}$. Moreover, we can also find a set $S \subseteq \mathcal{L}^\sigma$ such that S spans $\mathcal{L}^\sigma$, and S is an upper triangle square matrix with positive diagonal entries and non-positive entries elsewhere. Now, the vector $\nu_{\bar{\sigma}}$ is a solution to the relaxation $\mathcal{F}_{\mathcal{L}^\sigma}^{\mathbb{Z}}(\nu_{\bar{\sigma}})$ (non-negativity constraints are removed). Then, we can add appropriate non-negative multiples of the vectors in S to $\nu_{\bar{\sigma}}$ such that it becomes non-negative, and thus, we arrive at a feasible solution of $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$.

See Algorithm 4 for a description of the feasibility algorithm.

Finally, note that each individual step needed to compute a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ is performed during the project-and-lift algorithm 3. So, at the same time as computing a Markov basis, we can compute a feasible solution. Moreover, we can compute feasible solutions for many different fibers simultaneously, and thus, we can avoid repeating the same Gröbner basis computations for each fiber.

Example 6.2. *We apply the above method to find a feasible solution of equality constrained integer knapsack problems (see [2]):*

$$\mathcal{F} = \{x : Ax = b, x \in \mathbb{N}^n\}$$

where $A \in \mathbb{N}^{1 \times n}$ and $b \in \mathbb{N}$. Let $\mathcal{L} = \mathcal{L}_A := \{u : Au = \mathbf{0}, u \in \mathbb{Z}^n\}$. Then $\mathcal{F} = \mathcal{F}_{\mathcal{L}}(\nu)$ where $\nu \in \{x : Ax = b, x \in \mathbb{Z}^n\}$. Finding such a vector ν can be done in polynomial time using the HNF algorithm. If no such ν exists, then the original problem $\mathcal{F}$ is infeasible. Computing a Markov basis

Algorithm 4 Feasibility algorithm

Input: a lattice $\mathcal{L}$ and a vector $\nu \in \mathbb{Z}^n$.

Output: a feasible solution $x \in \mathcal{F}_{\mathcal{L}}(\nu)$ or infeasible.

Find a set $\sigma \subseteq \{1, \dots, n\}$ such that $\ker(\pi_{\sigma}) \cap \mathcal{L} = \{\mathbf{0}\}$.

Compute a set $M \subseteq \mathcal{L}^{\sigma}$ that is a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^{\sigma}$.

Compute a feasible solution $x \in \mathcal{F}_{\mathcal{L}^{\sigma}}(\nu_{\bar{\sigma}})$.

while $\sigma \neq \emptyset$ **do**

 Select $i \in \sigma$

if i is bounded **then**

$G := \mathcal{CP}_{\nu_{\bar{\sigma}}}(\prec_{-\omega^i}^{\sigma}, M)$

$M := (\pi_i^{\sigma})^{-1}(G)$

$x := \mathcal{NF}(x, G)$

$x := (\pi_i^{\sigma})^{-1}(x - \nu_{\bar{\sigma}}) + \pi_{\sigma \setminus i}(\nu)$

if $x_i < 0$ **then return** *infeasible*

else

 Compute $u \in \mathcal{L}^{\sigma \setminus i} \cap \mathbb{N}^n$ such that $u_i > 0$

$M := (\pi_i^{\sigma})^{-1}(M) \cup \{u\}$

$x := (\pi_i^{\sigma})^{-1}(x - \nu_{\bar{\sigma}}) + \pi_{\sigma \setminus i}(\nu)$

$x := x + \lambda u$ where $\lambda \in \mathbb{N}$ such that $(x + \lambda u)_i > 0$

end if

$\sigma := \sigma \setminus i$

$M := \{u \in M : u^+ \in \mathcal{B}_{\mathcal{L}^{\sigma}}(\nu_{\bar{\sigma}})\}$

end while

return x .

and thus solving the feasibility problem of any such knapsack problem involves only one Gröbner basis computation.²

Consider the following knapsack feasibility problem:

$$\mathcal{F} := \{x : 12223x_1 + 12224x_2 + 36674x_3 + 61119x_4 + 85569x_5 = 89643481 : x \in \mathbb{N}^5\}.$$

Let $\mathcal{L} = \mathcal{L}_A$ where $A = \begin{bmatrix} 12223 & 12224 & 36674 & 61119 & 85569 \end{bmatrix}$. Then, the set

$$S = \{(12224, -12223, 0, 0, 0), (2, -5, 1, 0, 0), (-1, -4, 0, 1, 0), (1, -8, 0, 0, 1)\}$$

spans $\mathcal{L}$. Let $\nu = (-4889, 12222, 0, 0, 0)$, then $\mathcal{F}_{\mathcal{L}}(\nu) = \mathcal{F}$. Let $\sigma = \{1\}$. Then, $\ker(\pi_{\sigma}) \cap \mathcal{L} = \{\mathbf{0}\}$, and also, we have $\nu_{\bar{\sigma}} = x = (12222, 0, 0, 0) \in \mathcal{F}_{\mathcal{L}^{\sigma}}(\nu_{\bar{\sigma}})$. A Markov basis of $\mathcal{L}^{\sigma}$ is

$$S' = \{(1, 0, 0, -1528), (0, 1, 0, -7640), (0, 0, 1, -6112), (0, 0, 0, 12223)\}.$$

Recall that $\omega^i \in \mathbb{Q}^{n-1}$ such that $\omega^i u = (\pi_i^{-1}(u))_i$. A $\prec_{-\omega^i}$ -Gröbner basis of $\mathcal{L}^{\sigma}$ is

$$G = \{(3, 2444, 0, 0), (-2, 2445, 0, 0), (5, -1, 0, 0), (-4, 0, 1, 0), (-3, -1, 0, 1)\}.$$

The normal form of x is $\mathcal{NF}(x, G) = x' = (2, 2444, 0, 0)$. We then lift this back into the original space, $x'' = (x' - \nu_{\bar{\sigma}}) + \nu = (-1, 2, 2444, 0, 0)$. This is not a feasible solution, and the problem is therefore infeasible.

²This Gröbner basis method for computing a feasible solution for the special case of equality constrained integer knapsack problems was found independently by Bjarke H. Roune ([15]).

We list the times to solve the feasibility problem in Figure 5. In each case, the problem was infeasible. The right hand sides used for each knapsack is the Frobenius number, that is, the largest infeasible right hand side.

Problem	Equality constraint (A)	RHS (b)	Time
cuww1	12223 12224 36674 61119 85569	89643481	0.00s
cuww2	12228 36679 36682 48908 61139 73365	89716838	0.00s
cuww3	12137 24269 36405 36407 48545 60683	58925134	0.00s
cuww4	13211 13212 39638 52844 66060 79268 92482	104723595	0.00s
cuww5	13429 26850 26855 40280 40281 53711 53714 67141	45094583	0.00s
prob1	25067 49300 49717 62124 87608 88025 113673 119169	3367335	0.00s
prob2	11948 23330 30635 44197 92754 123389 136951 140745	14215206	0.00s
prob3	39559 61679 79625 99658 133404 137071 159757 173977	58424799	0.02s
prob4	48709 55893 62177 65919 86271 87692 102881 109765	60575665	0.01s
prob5	28637 48198 80330 91980 102221 135518 165564 176049	62442884	0.01s
prob6	20601 40429 42407 45415 53725 61919 64470 69340 78539 95043	22382774	0.30s
prob7	18902 26720 34538 34868 49201 49531 65167 66800 84069 137179	27267751	0.00s
prob8	17035 45529 48317 48506 86120 100178 112464 115819 125128 129688	21733990	0.01s
prob9	13719 20289 29067 60517 64354 65633 76969 102024 106036 199930	13385099	0.01s
prob10	45276 70778 86911 92634 97839 125941 134269 141033 147279 153525	106925261	0.05s
prob11	11615 27638 32124 48384 53542 56230 73104 73884 112951 130204	577134	0.48s
prob12	14770 32480 75923 86053 85747 91772 101240 115403 137390 147371	944183	0.32s
prob13	15167 28569 36170 55419 70945 74926 95821 109046 121581 137695	6765260	0.78s
prob14	11828 14253 46209 52042 55987 72649 119704 129334 135589 138360	80230	0.23s
prob15	13128 37469 39391 41928 53433 59283 81669 95339 110593 131989	1663281	0.17s
prob16	35113 36869 46647 53560 81518 85287 102780 115459 146791 147097	109710	0.75s
prob17	14054 22184 29952 64696 92752 97364 118723 119355 122370 140050	752109	0.22s
prob18	20303 26239 33733 47223 55486 93776 119372 136158 136989 148851	783879	0.51s
prob19	20212 30662 31420 49259 49701 62688 74254 77244 139477 142101	677347	0.29s
prob20	32663 41286 44549 45674 95772 111887 117611 117763 141840 149740	1037608	0.45s

Figure 5: Hard Knapsack Constraint Instances.

In the paper [2], the feasibility is solved problem for the same set of equality constrained integer knapsack problems by using a reduced lattice basis approach. The solutions times in [2] and our solutions times are all less than a second, and so it would be interesting to compare the two methods on larger problems with a significant computation time.

This approach for computing a feasible solution of a fiber could potentially be used when computing a truncated Markov basis by the project and lift algorithm since during the algorithm, we check whether $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu)$ which is the feasibility problem $\mathcal{F}_{\mathcal{L}}(\nu - \nu') \neq \emptyset$. Note that the feasibility approach is well-suited to computing feasibility for many different fibers simultaneously. It would be interesting to see the performance of this approach.

7 Optimality

In this section, we discuss the use of Gröbner bases to solve the integer program

$$IP_{\mathcal{L},c}(\nu) := \min\{cx : x \in \mathcal{F}_{\mathcal{L}}(\nu)\}.$$

The most straight-forward way to solve $IP_{\mathcal{L},c}(\nu)$ is to compute a Markov basis of $\mathcal{L}$ and a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ and then compute a $\prec_c$ -Gröbner basis of $\mathcal{L}$ for some term order $\prec$, and finally, compute the normal form of the feasible solution giving the optimal solution. Here, we are actually solving $IP_{\mathcal{L},\prec_c}(\nu)$ which is essentially the same as solving $IP_{\mathcal{L},c}(\nu)$. With this method, if we want to solve $IP_{\mathcal{L},c}(\nu')$ for a finite number of ν' , we only to compute a feasible solution of $\mathcal{F}_{\mathcal{L}}(\nu)$ and redo the normal form computation without needing to recompute the Markov basis or the Gröbner basis. Also, note that the feasible solutions can be computed at the same time as computing the Markov basis without much additional computational overhead (see Section 6).

If we wish to solve $IP_{\mathcal{L},c}(\nu)$ for just one ν , then we should use information specific to that fiber to solve the problem. We can compute a ν -truncated Markov basis $\mathcal{L}$ and a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$. The problem with this method is that we must compute the entire ν -truncated $\prec_c$ -Gröbner basis in order to prove optimality. Moreover, if the feasible set $\mathcal{F}_{\mathcal{L}}(\nu)$ is large, then the truncated Gröbner basis may still be quite large and in some cases as large as the non-truncated Gröbner basis. Thus, we need further ways of reducing its size.

Some of the non-negativity constraints on the x variables may not strictly be necessary to solve the problem since they may be redundant or not active near the optimal solution. So, we consider relaxations of $IP_{\mathcal{L},c}(\nu)$ in which we relax the non-negativity constraints on some of the x variables. Consider the problem

$$IP_{\mathcal{L},c}^{\sigma}(\nu) := \min\{cx : x \equiv \nu \pmod{\mathcal{L}}, x_{\bar{\sigma}} \geq 0, x \in \mathbb{Z}^n\}$$

where $\sigma \subseteq \{1, \dots, n\}$. Here, we have relaxed the non-negativity constraints on x_{σ} . If σ is the set of basic variables given by solving the linear relaxation of $IP_{\mathcal{L},c}(\nu)$ using the simplex algorithm (see [14]), the relaxation $IP_{\mathcal{L},c}^{\sigma}(\nu)$ is called a *group relaxation* (see [9]). Note that, for group relaxations, $IP_{\mathcal{L},c}^{\sigma}(\nu)$ has an optimal solution and $\ker(\pi_{\sigma}) \cap \mathcal{L} = \{\mathbf{0}\}$. If σ is any subset of the above set for the group relaxation, then $IP_{\mathcal{L},c}^{\sigma}(\nu)$ is called an *extended group relaxation* (see [23]). Thus, the original problem $IP_{\mathcal{L},c}(\nu) = IP_{\mathcal{L},c}^{\emptyset}(\nu)$ is an extended group relaxation.

We want to solve these extended group relaxations using Gröbner bases and so we must rewrite $IP_{\mathcal{L},c}^{\sigma}(\nu)$ in the form $IP_{\mathcal{L}',c'}^{\sigma}(\nu')$ for some lattice $\mathcal{L}'$, some right hand side ν' , and some cost function c' . Firstly, any extended group relaxation $IP_{\mathcal{L},c}^{\sigma}(\nu)$ that has an optimal solution can always be rewritten in the equivalent form $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$ for some $\tilde{c}$ where $\tilde{c}_{\sigma} = \mathbf{0}$, that is, $\tilde{c}_i = 0$ for all $i \in \sigma$ (as given by the simplex algorithm [14]) where $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$ has the same optimal solution as $IP_{\mathcal{L},c}^{\sigma}(\nu)$, although the optimal value may differ by a known constant. Now consider the projection of $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$ onto the $\bar{\sigma}$ components:

$$IP_{\mathcal{L}^{\sigma},\tilde{c}_{\bar{\sigma}}}(\nu_{\bar{\sigma}}) := \min\{\tilde{c}_{\bar{\sigma}}x : x \equiv \nu_{\bar{\sigma}} \pmod{\mathcal{L}^{\sigma}}, x \in \mathbb{N}^{|\bar{\sigma}|}\}.$$

Any feasible solution of $IP_{\mathcal{L}^{\sigma},\tilde{c}_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ lifts to a feasible solution of $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$. Let $x \in \mathcal{F}_{\mathcal{L}^{\sigma}}(\nu_{\bar{\sigma}})$, then $x' := (x - \pi_{\bar{\sigma}}(\nu)) + \nu \in \mathcal{F}_{\mathcal{L}}^{\sigma}(\nu)$. Moreover, an optimal solution of $IP_{\mathcal{L}^{\sigma},\tilde{c}_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ lifts to an optimal solution of $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$ since $\tilde{c}_{\bar{\sigma}}x = \tilde{c}x'$. Hence, $IP_{\mathcal{L},c}^{\sigma}(\nu)$, $IP_{\mathcal{L},\tilde{c}}^{\sigma}(\nu)$ and $IP_{\mathcal{L}^{\sigma},\tilde{c}_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ are all essentially equivalent problems.

The basic idea of the algorithm is that we start with the group relaxation $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ where σ is defined as above and we assume $c_{\bar{\sigma}} = \mathbf{0}$ without loss of generality. We then solve the group relaxation $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$. If the optimal solution of $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ lifts to a feasible solution of $IP_{\mathcal{L}, c}(\nu)$, then it is optimal for $IP_{\mathcal{L}, c}(\nu)$ and we are done. Otherwise, we add a non-negativity constraint on one of the unconstrained x variables, that is, we choose $i \in \sigma$ and set $\sigma := \sigma \setminus i$, and solve the extended group relaxation $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$. Again if the optimal solution of $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ lifts to a feasible for $IP_{\mathcal{L}, c}(\nu)$, then we are done. Otherwise, we again add a non-negativity constraint on one of the unconstrained x variables and solve the new extended group relaxation, and so on, until $IP_{\mathcal{L}, c}(\nu)$ is solved. The algorithm must terminate with a solution because in the worst case we end up solving the original problem $IP_{\mathcal{L}, c}(\nu)$ ($\sigma = \emptyset$).

To solve an extended group relaxation $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$, we first compute a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^\sigma$ and a feasible solution of $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$. Secondly, we compute a $\nu_{\bar{\sigma}}$ -truncated $\prec_{c_{\bar{\sigma}}}^\sigma$ -Gröbner basis of $\mathcal{L}^\sigma$. Then, we compute the normal form of the feasible solution giving the optimal solution of $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$.

Initially, we need to compute a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^\sigma$ for the group problem. Let B be a basis of the lattice $\mathcal{L}$. Then for the group problem, σ gives $\text{rank}(B)$ linear independent columns of B . Thus, as discussed at the end of Section 5, we can compute a Markov basis via a HNF computation. Similarly, we can compute a feasible solution as discussed at the end of Section 6.

At each iteration of the algorithm, we could compute a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^\sigma$ starting from scratch each time as described in Section 5, but instead, we can compute it incrementally exactly as in the Project-and-Lift algorithm. In the previous iteration, we will have computed a $\nu_{\bar{\sigma}'}$ -truncated Markov basis of $\mathcal{L}^{\bar{\sigma}'}$ where $\bar{\sigma}' := \sigma \cup i$ for some $i \in \bar{\sigma}$. Hence, using Lemmas 5.1 and 5.3, we can compute a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^\sigma$ from a $\nu_{\bar{\sigma}'}$ -truncated Markov basis of $\mathcal{L}^{\bar{\sigma}'}$ in one step. In effect, we perform the Project-and-Lift algorithm simultaneously. Also, as discussed in Section 6, we can compute a feasible solution of $IP_{\mathcal{L}^\sigma, c_{\bar{\sigma}}}(\nu_{\bar{\sigma}})$ from a feasible solution of $IP_{\mathcal{L}^{\bar{\sigma}'}, c_{\bar{\sigma}'}}(\nu_{\bar{\sigma}'})$.

See Algorithm 5 for a description of the optimisation algorithm.

Algorithm 5 Optimisation algorithm

Input: an integer program $IP_{\mathcal{L}, c}(\nu)$.

Output: an optimal solution $x \in \mathcal{F}_{\mathcal{L}}(\nu)$ or *infeasible*.

Find a set $\sigma \subseteq \{1, \dots, n\}$ such that $\ker(\pi_\sigma) \cap \mathcal{L} = \{\mathbf{0}\}$.

while $\sigma \neq \emptyset$ **do**

 Compute a set $M \subseteq \mathcal{L}^\sigma$ that is a $\nu_{\bar{\sigma}}$ -truncated Markov basis of $\mathcal{L}^\sigma$.

 Compute a feasible solution $x \in \mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}})$ and **return** *infeasible* if $\mathcal{F}_{\mathcal{L}^\sigma}(\nu_{\bar{\sigma}}) = \emptyset$.

$G := \mathcal{CP}_{\nu_{\bar{\sigma}}}(\prec_{c_{\bar{\sigma}}}^\sigma, M)$

$x := \mathcal{NF}(x, G)$

$x := \pi_\sigma^{-1}(x - \nu_{\bar{\sigma}}) + \nu$

if $x \geq 0$ **then return** x

 Select $i \in \sigma$.

$\sigma := \sigma \setminus i$

end while

At each iteration, we must select the next i . An obvious choice is to select the component with the most violated non-negativity constraint, that is, the most negative component.

Existing branch-and-bound methods for integer programming can take advantage of a good feasible solution of $IP_{\mathcal{L},c}(\nu)$, but the above method cannot. However, we can take advantage of a good feasible solution since a feasible solution gives us an upper bound on $IP_{\mathcal{L},c}(\nu)$ that can be used to strengthen truncation.

Consider the following reformulation of $IP_{\mathcal{L},c}(\nu)$ using some upper bound $k \in \mathbb{Z}$ on $IP_{\mathcal{L},c}(\nu)$:

$$\begin{aligned} IP_{\mathcal{L},c}(\nu) &:= \min\{cx : x \equiv \nu \pmod{\mathcal{L}}, x \in \mathbb{N}^n\} \\ &= \min\{cx : x \equiv \nu \pmod{\mathcal{L}}, cx \leq k, x \in \mathbb{N}^n\} \\ &= \min\{-y : x \equiv \nu \pmod{\mathcal{L}}, cx + y = k, x \in \mathbb{N}^n, y \in \mathbb{N}\} + k. \end{aligned}$$

To solve the reformulation, we first need to express it in the form $IP_{\mathcal{L}',c'}(\nu')$ for some lattice $\mathcal{L}'$, right hand side ν' and some cost function c' . Let $\mathcal{L}' := \{(u, -cu) : u \in \mathcal{L}\}$, $\nu' := (\nu, k - c\nu)$, and $c' = (\mathbf{0}, -1)$. Then,

$$\begin{aligned} IP_{\mathcal{L}',c'}(\nu') &:= \min\{-y : (x, y) \equiv (\nu, k - c\nu) \pmod{\mathcal{L}'}, (x, y) \in \mathbb{N}^{n+1}\} \\ &= \min\{-y : (x - \nu, y - (k - c\nu)) \in \mathcal{L}', (x, y) \in \mathbb{N}^{n+1}\} \\ &= \min\{-y : (x - \nu, y - (k - c\nu)) = (u, -cu), u \in \mathcal{L}, (x, y) \in \mathbb{N}^{n+1}\} \\ &= \min\{-y : x \equiv \nu \pmod{\mathcal{L}}, cx + y = k, x \in \mathbb{N}^n, y \in \mathbb{N}\}. \end{aligned}$$

Thus, $IP_{\mathcal{L},c}(\nu) = IP_{\mathcal{L}',c'}(\nu') + k$. So, we can solve $IP_{\mathcal{L},c}(\nu)$ by solving $IP_{\mathcal{L}',c'}(\nu')$ using the methods discussed previously.

Hopefully, a ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$ is a lot smaller than a ν -truncated Gröbner basis of $\mathcal{L}$. Crucially, the size of a minimal ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$ cannot exceed the size of a minimal ν -truncated $\prec_c$ -Gröbner basis, and so, we are not computing more than before. This follows from Lemma 5.1; since $\pi_{n+1}(\mathcal{L}') = \mathcal{L}$, $\pi_{n+1}(\nu') = \nu$, $c' = -e^{n+1}$, and $c = -\omega^{n+1}$, if G is a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$, then $\pi_{n+1}^{-1}(G)$ is a ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$. How much smaller a ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$ is than a ν -truncated Gröbner basis of $\mathcal{L}$ will depend on the strength of the upper bound. It is potentially just the empty set.

If we are given a feasible solution $x \in \mathcal{F}_{\mathcal{L}}(\nu)$, then we can set $k = cx - 1$. If x is the optimal solution of $IP_{\mathcal{L},c}(\nu)$, then setting $k = cx - 1$, we have $\mathcal{F}_{\mathcal{L}'}(\nu') = \emptyset$. Therefore, a minimal ν' -truncated Markov basis of $\mathcal{L}'$ is empty and a minimal ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$ is also empty. So, potentially, computing a ν' -truncated $\prec_{c'}$ -Gröbner basis of $\mathcal{L}'$ is a lot more efficient than computing a ν -truncated $\prec_c$ -Gröbner basis of $\mathcal{L}$; we only have to compute an empty set to show optimality!

Even if the bound k is not very good and thus does not help truncation much, it is still definitely worthwhile to solve $IP_{\mathcal{L}',c'}(\nu')$ instead of $IP_{\mathcal{L},c}(\nu)$. The reason is that by introducing the constraint $cx \leq k$ into the problem, more components may become *bounded* and thus the Gröbner basis and the Markov basis computations for $IP_{\mathcal{L}',c'}(\nu')$ are faster (see the section in [12] on Criterion 2). The computations are also faster for the extended group relaxations of $IP_{\mathcal{L}',c'}(\nu')$ as well.

Example 7.1. Let $\mathcal{L} = \mathcal{L}_A$ for the matrix A given in Example 4.7. In the following table, we list the time taken to compute the optimal solution of $IP_{\mathcal{L},c}(\nu)$ for different ν 's given a feasible solution where $c = (3, 15, 1, 5, 2, 17, 16, 16, 15, 9, 7, 11, 13)$.

ν	<i>Group Relaxation</i>	<i>Final Relaxation</i>	<i>Time</i>
(1, 1, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 0)	{3, 4, 5, 10}	{3, 4, 5, 10}	0.28s
(1, 0, 1, 0, 3, 0, 1, 5, 0, 1, 0, 9, 0)	{1, 5, 8, 12}	{1, 5, 8, 12}	0.02s
(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)	{3, 4, 5, 10}	{5, 10}	0.61s
(1, 2, 0, 3, 5, 0, 1, 3, 0, 4, 0, 1, 0)	{3, 4, 5, 10}	{3, 5, 10}	0.46s
(19, 7, 3, 8, 13, 11, 1, 15, 4, 8, 17, 9, 5)	{3, 4, 5, 10}	{3, 4, 5, 10}	0.28s

Here, we used the quick check for truncation.

If we do not have a good feasible solution or any feasible solution at all available, we can still use the extended formulation. Assume that we are given a lower bound l on the optimal value, which we can always find by solving the linear relaxation. We then try to find a feasible solution by computing a ν' -truncated Markov basis of $\mathcal{L}'$ where $k = l$. If we find a feasible solution, then it must be optimal. Otherwise, we recompute a ν' -truncated Markov basis of $\mathcal{L}'$ where $k = l + 1$ and again try to find a feasible solution. We repeat this procedure by incrementing k until we find a feasible solution which must be an optimal solution. This procedure has the advantage that we only compute feasible solutions and not optimal solutions, and thus, we avoid some Gröbner basis computations.

The recomputation of ν' -truncated Markov bases to find initial solutions initially seems inefficient; however, this is not the case because we can reuse the previous computations. Let $\nu' = (\nu, k - c\nu)$ and $\nu'' = (\nu, k + 1 - c\nu)$. Then $\nu' \in \mathcal{B}_{\mathcal{L}}(\nu'')$ since $(\mathbf{0}, 1) \in \mathcal{F}_{\mathcal{L}'}(\nu'' - \nu')$ and so $\mathcal{B}_{\mathcal{L}}(\nu') \subseteq \mathcal{B}_{\mathcal{L}}(\nu'')$. Hence, to compute a ν'' -truncated Gröbner basis or Markov basis of $\mathcal{L}$ requires also computing a ν' -truncated Gröbner basis or Markov basis respectively anyway. This applies not only at the final stage of the algorithm for each value of k , but also at each intermediate stage for the extended group relaxations of $IP_{\mathcal{L}', c'}(\nu')$. So, it requires keeping all the intermediate stages around, and is thus more complex than the first algorithm presented in this section.

We have not implemented such an approach yet. It would be interesting to see how it performs. However, if we know a very good initial feasible solution, we would expect that the previous method is faster.

8 Conclusion

We have demonstrated that it is possible to significantly improve upon the performance of previous Gröbner basis based approaches. However, Gröbner basis approaches have not yet proven to be competitive with traditional branch-and-bound based approaches to integer programming for industrial applications, but given the significant advances shown here, perhaps with further research, Gröbner basis methods will be useful for some classes of problems when combined with existing methods.

A Lattice Programs and Integer Programs

In this appendix, we show the equivalence of fibers and feasible sets of integer programs and of lattice programs and integer programs.

Consider the set $\{x : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\}$ where $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Z}^m$, $\sigma \subseteq \{1, \dots, n\}$, $\bar{\sigma}$ is the complement of σ , and $x_{\bar{\sigma}}$ is the set of variables indexed by $\bar{\sigma}$. Thus, the $x_{\bar{\sigma}}$ variables are non-negative and the x_{σ} variables (the x variables indexed by σ) are unrestricted in sign. The set of feasible solutions for any integer program can be represented in this form. We will rewrite this set as a lattice fiber. We will actually only consider the projection of this set onto the non-negative variables (the $\bar{\sigma}$ components), that is, $F_A^{\sigma}(b) := \{x_{\bar{\sigma}} : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\}$ since any solution of $F_A^{\sigma}(b)$ can easily be extended to a solution in the original space; more specifically, given $\tilde{x}_{\bar{\sigma}} \in F_A^{\sigma}(b)$, we can find $\tilde{x}_{\sigma} \in \mathbb{Z}^{|\sigma|}$ where $A_{\sigma}\tilde{x}_{\sigma} = b - A_{\bar{\sigma}}\tilde{x}_{\bar{\sigma}}$ by using the HNF algorithm where A_{σ} and $A_{\bar{\sigma}}$ are the sub-matrices of A whose columns are indexed by σ and $\bar{\sigma}$ respectively.

We now write $F_A^{\sigma}(b)$ as a fiber of a lattice. Let $\mathcal{L}_A = \{u \in \mathbb{Z}^n : Au = \mathbf{0}\}$, and let $\nu \in \mathbb{Z}^n$ where $A\nu = b$. Note that we can compute a basis of $\mathcal{L}_A$ and a ν using the HNF algorithm. Also, let $\mathcal{L}_A^{\sigma}$ be the lattice $\mathcal{L}_A$ projected onto the $\bar{\sigma}$ components (i.e. $\mathcal{L}_A^{\sigma} := \{u_{\bar{\sigma}} : Au = \mathbf{0}\}$). Note that a projection of a lattice is always again a lattice. It follows that

$$\begin{aligned} F_A^{\sigma}(b) &= \{x_{\bar{\sigma}} : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} \\ &= \{x_{\bar{\sigma}} : Ax = A\nu, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} \\ &= \{x_{\bar{\sigma}} : A(x - \nu) = \mathbf{0}, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} \\ &= \{x_{\bar{\sigma}} : x - \nu \in \mathcal{L}_A, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} \\ &= \{x_{\bar{\sigma}} : x_{\bar{\sigma}} - \nu_{\bar{\sigma}} \in \mathcal{L}_A^{\sigma}, x_{\bar{\sigma}} \geq \mathbf{0}, x_{\bar{\sigma}} \in \mathbb{Z}^{|\bar{\sigma}|}\} \\ &= \mathcal{F}_{\mathcal{L}_A^{\sigma}}(\nu_{\bar{\sigma}}). \end{aligned}$$

So, the set of feasible solutions to any integer program can be represented as a fiber of a lattice.

Conversely, given a $\mathcal{L}$ and a $\nu \in \mathbb{Z}^n$, we can represent the fiber $\mathcal{F}_{\mathcal{L}}(\nu)$ as the set of feasible solutions to an integer program. Let $S \in \mathbb{Z}^{n \times k}$ be a matrix where the columns of S span the lattice $\mathcal{L}$. Then,

$$\begin{aligned} \mathcal{F}_{\mathcal{L}}(\nu) &= \{x \in \mathbb{Z}^n : x \equiv \nu \pmod{\mathcal{L}}\} \\ &= \{x \in \mathbb{Z}^n : x - \nu \in \mathcal{L}\} \\ &= \{x \in \mathbb{Z}^n : x - \nu = Sy, y \in \mathbb{Z}^k\} \\ &= \{x : (I, -S)(x, y) = \nu, x \geq 0, x \in \mathbb{Z}^n, y \in \mathbb{Z}^k\}. \\ &= \{x : A(x, y) = \nu, x \geq 0, x \in \mathbb{Z}^n, y \in \mathbb{Z}^k\}. \\ &= F_A^{\sigma}(\nu) \end{aligned}$$

where $A = (I, -S) \in \mathbb{Z}^{(n+k) \times n}$ and $\sigma = \{1, \dots, n\}$ and $\bar{\sigma} = \{n+1, \dots, n+k\}$. Thus, fibers of lattices and feasible sets of integer programs are two different representations of the same set and so are equivalent concepts.

Next we show the equivalence of lattice programs and integer programs. Any integer linear program can be written in the form $\min\{cx : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\}$ where $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Z}^m$, and $c \in \mathbb{Z}^n$. If this integer program has an optimal solution, then there exists a $\tilde{c} \in \mathbb{Z}^n$ where $\tilde{c}_{\sigma} = \mathbf{0}$ such that $cx = \tilde{c}x + k$ for some constant $k \in \mathbb{Q}^n$ and every feasible solution x . Then,

$$\begin{aligned} \min\{cx : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} &= \min\{\tilde{c}x + k : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} \\ &= \min\{\tilde{c}_{\bar{\sigma}}x_{\bar{\sigma}} : Ax = b, x_{\bar{\sigma}} \geq \mathbf{0}, x \in \mathbb{Z}^n\} + k \\ &= \min\{\tilde{c}_{\bar{\sigma}}x_{\bar{\sigma}} : x_{\bar{\sigma}} \in F_A^{\sigma}(b)\} + k \\ &= \min\{\tilde{c}_{\bar{\sigma}}x_{\bar{\sigma}} : x_{\bar{\sigma}} \in \mathcal{F}_{\mathcal{L}_A^{\sigma}}(\nu_{\bar{\sigma}})\} + k \end{aligned}$$

where $\mathcal{L}_A^\sigma$ and $\nu_{\bar{\sigma}}$ are defined as above. More specifically, we can always find a $\tilde{c} \in \mathbb{Z}^n$ where $\tilde{c}_{\bar{\sigma}} = \mathbf{0}$ such that $cx = \tilde{c}x + k$ for every feasible solution x if and only if $cx = 0$ for all $x \in \mathbb{Z}^n$ where $x_{\bar{\sigma}} = \mathbf{0}$ and $Ax = \mathbf{0}$. Note that if $cx \neq 0$ for some $x \in \mathbb{Z}^n$ where $x_{\bar{\sigma}} = \mathbf{0}$ and $Ax = \mathbf{0}$, then the integer program has no optimal solution. We can check this condition and find a valid $\tilde{c}$ easily using the HNF algorithm.

Hence, any integer program with an optimal solution can be written in the form

$$IP_{\mathcal{L},c}(\nu) := \min\{cx : x \in \mathcal{F}_{\mathcal{L}}(\nu)\}$$

for some lattice $\mathcal{L} \subseteq \mathbb{Z}^n$, a vector $\nu \in \mathbb{Z}^n$, and a vector $c \in \mathbb{Z}^n$. As we saw at the end of Section 3, solving $IP_{\mathcal{L},c}(\nu)$ is equivalent to solving the *lattice program* $IP_{\mathcal{L},\succ}(\nu)$ where $\succ$ is compatible with c .

Acknowledgments

I would like to thank Raymond Hemmecke and Laurence Wolsey for many fruitful discussions and helpful comments.

References

- [1] 4ti2 team. 4ti2 – a software package for algebraic, geometric and combinatorial problems on linear spaces. Available at www.4ti2.de.
- [2] K. Aardal and A. K. Lenstra. Hard equality constrained integer knapsacks. *Mathematics of operations research*, 29(3):724–738, 2004.
- [3] A.M. Bigatti, R. LaScala, and L. Robbiano. Computing toric ideals. *Journal of Symbolic Computation*, 27:351–365, 1999.
- [4] B. Buchberger. History and basic features of the critical-pair/completion procedure. *Journal of Symbolic Computation*, 2:3–38, 1987.
- [5] P. Conti and C. Traverso. Buchberger algorithm and integer programming. In *Proceedings AAECC-9 (New Orleans)*, volume 539 of *LNCS*, pages 130–139. Springer Verlag, 1991.
- [6] D. Cox, J. Little, and D. O’Shea. *Ideals, Varieties and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Springer Verlag, 1992.
- [7] P. Diaconis and B. Sturmfels. Algebraic algorithms for sampling from conditional distributions. *Annals of Statistics*, 26:363–397, 1998.
- [8] N. Eriksson. Toric ideals of homogeneous phylogenetic models. In *Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation*, pages 149–154. ACM Press, 2004.
- [9] Ralph E. Gomory. On the relation between integer and noninteger solutions to linear programs. In *Proceedings of the National Academy of Sciences*, volume 53, pages 260–265, 1965.

- [10] Jack E. Graver. On the foundation of linear and integer linear programming I. *Mathematical Programming*, 9:207–226, 1975.
- [11] R. Hemmecke. On the computation of Hilbert bases and extreme rays of cones. e-print: arXiv:math.CO/0203105, 2002.
- [12] R. Hemmecke and Peter N. Malkin. Computing generating sets of lattice ideals. e-print: arXiv:math.CO/0508359, 2006.
- [13] Serkan Hosten and Bernd Sturmfels. GRIN: An implementation of Gröbner bases for integer programming. In E. Balas and J. Clausen, editors, *Integer Programming and Combinatorial Optimization*, volume 920 of *LNCS*, pages 267–276. Springer Verlag, 1995.
- [14] George L. Nemhauser and Laurence A. Wolsey. *Integer and Combinatorial Optimization*. John Wiley and Sons, New York, 1988.
- [15] Bjarke H. Roune. A gröbner basis attack on the frobenius problem. private communication, 2006.
- [16] Bernd Sturmfels. *Gröbner bases and convex polytopes*, volume 8 of *University Lecture Notes Series*. American Mathematical Society, Providence, Rhode Island, 1996.
- [17] Rekha R. Thomas. The structure of group relaxations. In Karen Aardal, George Nemhauser, and Robert Weismantel, editors, *Discrete Optimization*, volume 12 of *Handbooks in Operations Research and Management Science*. Elsevier, 2005.
- [18] Rekha Thomas and Robert Weismantel. Truncated gröbner bases for integer programming. *Applicable Algebra in Engineering, Communication and Computing*, 8:241–257, 1997.
- [19] Rekha R. Thomas. A geometric Buchberger algorithm for integer programming. *Mathematics of Operations Research*, 20:864–884, 1995.
- [20] Rekha R. Thomas and Serkan Hosten. Gomory integer programs. *Mathematical Programming Series B*, 96:271–292, 2003.
- [21] R. Urbaniak, R. Weismantel, and Günter M. Ziegler. A variant of Buchberger’s algorithm for integer programming. *SIAM J. on Discrete Mathematics*, 10:96–108, 1997.
- [22] R. Weismantel. Test sets of integer programs. *Mathematical Methods of Operations Research*, 47:1–37, 1998.
- [23] L. Wolsey. Extensions of the group theoretic approach in integer programming. *Management Science*, 18(1):74–83, 1971.